Hip and Knee Orthopaedic clinic

Outpatient service information, triage and referral guidelines

Clinical features indicative of potentially serious pathology

General hip and knee

Hip fracture - hip pain associated with history of trauma

Malignancy - hip or knee symptoms associated with red flags symptoms

- Night pain
- Anorexia/weight loss
- Malaise +/- history of cancer

Septic joint – severe pain, joint swelling and systematically unwell patient

Peri-prosthetic fractures - severe pain, inability to mobilise

Post-operative patients

Elective Surgery

Symptoms - severe pain, swelling increased local temperature, history of trauma / injury to relevant site post-surgery

Infection - wound related issues, cellulitis, swelling, systematically unwell patient

Prosthetic hip dislocation

DVT/PE – calf swelling, pain, tender to touch – For GP investigation and if positive discuss with orthopaedic registrar on-call via the RAH switchboard on (08) 7074 0000 or TQEH switchboard (08) 8222 6000 regarding anti-coagulation trauma.

Compartment syndrome - pain excessive of that expected for injury, swelling

Infection – wound related issues, cellulitis, swelling, systematically unwell patient.

General information to assist with referrals and the referral templates for the RAH & TQEH are available to download from the CALHN Outpatient Services website

For more information or to make a referral

Royal Adelaide Hospital: Level 3G, Wing 2 Referral Fax Number: (08) 7074 6247 Phone Number: 1300 153 853

The Queen Elizabeth Hospital: Ground Floor, Area 3

Referral Fax Number: (08) 8222 6233 Phone Number: (08) 8222 7020

Web: www.sahealth.gov.au

Acknowledgement: This document has had the final approval from the head of unit:

Dr ???????

Version	Approved date	Review date	Ammendment
Draft V0.1	20181005	20191005	



