



Rapid Detection and Response Paediatric Observation Chart
(0 - 3 months)
MR-59B

Affix patient identification label in this box

U.R. No:.....

Surname:.....

Given Name:.....

Second Given Name:.....

D.O.B.: Sex/Gender:

Chart Number: Mid Arm circumference: Height: Weight:

SECTION A - GENERAL INSTRUCTIONS

Minimum set of observations – Write in Section C
Take observations on child (at rest and record) on admission:

- Respiratory rate, oxygen saturation SpO₂, blood pressure, pulse rate, temperature, pain score, level of consciousness
- Other observations as indicated including BGL, O₂ Flow rate, O₂ delivery method, capillary refill and level of sedation

How to record observations in Section C
Place a dot (.) in the centre of the box that includes the current observation in its range of values. Connect the new dot to the previous dot with a straight line. Write the value in the relevant box for O₂ flow rate, BGL, and also if observations fall above or below graphic parameters as indicated.

For systolic blood pressure use the symbol indicated on the graphic chart. Use the right arm (unless contraindicated) to measure blood pressure. Document cuff size and the 95th percentile for this baby/child (at Section C). Refer to Section D (Modifications) for the blood pressure limits that trigger MDT review for this baby/child.

Other Observations
Level of consciousness should be documented using the AVPU scale except for children receiving sedation and/or opioids, where a level of sedation score should be recorded in place of the level of consciousness.
Select pain assessment tool appropriate for the age, developmental level and clinical state of the child. Refer to state and/or local guidelines for pain assessment tools.

SECTION B - ASSESSMENT OF RESPIRATORY DISTRESS

Used together with Respiratory Rate to provide further information about the airway and breathing assessment. Not all features may be present. Escalate as indicated.

| | MILD | MODERATE | SEVERE |
|----------------------------|---|--|---|
| Airway | Stridor only with exertion / crying | Some stridor at rest | Biphasic or increasing severity of stridor at rest |
| Work of breathing | Mild chest retraction (intercostal and/or suprasternal recession) | Moderate chest retraction (moderate intercostal and/or suprasternal recession) Tracheal tug / head bob / nasal flaring may be present | Severe chest retraction (marked intercostal, suprasternal and sternal recession) Tracheal tug / head bob / nasal flaring Grunting / gasping |
| Colour | Pink | Pallor | Dusky, mottled, cyanotic, extreme pallor |
| Behaviour / feeding | Normal behaviour / interactive No difficulty feeding Talks in sentences Loud cry | Intermittent irritability / difficult to console / more tired than usual Difficulty feeding Some difficulty talking (words only) | Agitated / confused or lethargic / looks exhausted Refuses / unable to feed Unable to talk or cry (too breathless) |
| Apnoea | Transient No desaturation | Transient with brief desaturations | Apnoea that is recurrent or prolonged or requires intervention |
| Oxygen | No oxygen requirement | New or increasing oxygen requirement | Hypoxaemia (SpO ₂ < 90% on Oxygen, HHHFNO or CPAP) |

RDR Paediatric Observation Chart (0-3 months) MR-59B

Rapid Detection and Response Paediatric Observation Chart
(0 - 3 months)
MR-59B

Affix patient identification label in this box

U.R. No:.....

Surname:.....

Given Name:.....

Second Given Name:.....

D.O.B.: Sex/Gender:

SECTION G - RESPONSE CRITERIA AND ACTIONS TO TAKE

ALWAYS CHECK CURRENT MODIFICATIONS

MEDICAL EMERGENCY RESPONSE (MER) CALL

| RESPONSE CRITERIA - If one or more observations are in the purple zone, or one or more of the following are occurring; | | ACTIONS REQUIRED |
|---|---|--|
| <ul style="list-style-type: none"> You are worried about the patient A patient or consumer is worried | <ul style="list-style-type: none"> Respiratory or cardiac arrest Threatened airway Significant bleeding Unexpected or uncontrolled seizure Consider for delayed MDT review (> 30 minutes) | <ul style="list-style-type: none"> Place emergency call and specify location Initiate basic/advanced life support Notify senior doctor responsible for patient Increase frequency of observations post intervention. Take advice from MER team |

MULTI DISCIPLINARY TEAM (MDT) REVIEW (Minimum team of registered nurse/midwife and medical practitioner)

| RESPONSE CRITERIA - If one or more observations are in the red zone, or one or more of the following are occurring; | | ACTIONS REQUIRED |
|---|--|---|
| <ul style="list-style-type: none"> You are worried about the patient A patient or consumer is worried | <ul style="list-style-type: none"> Poor peripheral circulation Greater than expected fluid loss Urine output < 1ml/kg/hr over 4 hours or patient has not voided for 8 hours New or increase in O₂ flow rate Escalate to MER call if there are 3 or more observations in red zone | <ul style="list-style-type: none"> MDT review must occur within 30 minutes (Rural Hospitals refer to local guidelines) or escalate to MER call Increase frequency of observations (minimum hourly) Escalate if there are ongoing fluctuations. Review SpO₂ and O₂ flow rate requirements |


REGISTERED NURSE OR REGISTERED MIDWIFE (and notify Shift Coordinator)

| RESPONSE CRITERIA - If one or more observations are in the yellow zone, or one or more of the following are occurring; | | ACTIONS REQUIRED |
|---|---|---|
| <ul style="list-style-type: none"> You are worried about the patient A patient or consumer is worried | <ul style="list-style-type: none"> Poor peripheral circulation New or unexplained behavioural change Unrelieved or unexpected pain Escalate to MDT review if there are 3 or more observations in yellow zone | <ul style="list-style-type: none"> Registered nurse/midwife review must occur within 30 minutes, or escalate to MDT review Increase frequency of observations Manage anxiety, pain and other symptoms Review SpO₂ and O₂ flow rate requirements |

SECTION H - SEDATION SCORE

| Score | Descriptor | Stimulus | Response | Duration |
|-------|---|------------------------|--|--------------|
| 3 | Difficult to rouse | Pain, shoulder squeeze | Brief eye opening OR any movement OR no response | N/A |
| 2 | Easy to rouse, difficulty staying awake | Voice, light touch | Eye opening and eye contact | < 10 seconds |
| 1 | Easy to rouse | Voice, light touch | Eye opening and eye contact | ≥ 10 seconds |
| 0 | Awake, alert when approached | N/A | N/A | N/A |

| SECTION C - OBSERVATION CHART | | | | | | | | | | | | |
|---|---------------------------|---------------|--|--|--|--|--|--|--|----------------|---------------------|---------------|
| Date | | | | | | | | | | | | |
| Time | | | | | | | | | | | | |
| Respiratory Rate (breaths/min) | Write ≥ 80 | | | | | | | | | | Write ≥ 80 | |
| | 75 - 79 | | | | | | | | | | 75 - 79 | |
| | 70 - 74 | | | | | | | | | | 70 - 74 | |
| | 65 - 69 | | | | | | | | | | 65 - 69 | |
| | 60 - 64 | | | | | | | | | | 60 - 64 | |
| | 55 - 59 | | | | | | | | | | 55 - 59 | |
| | 50 - 54 | | | | | | | | | | 50 - 54 | |
| | 45 - 49 | | | | | | | | | | 45 - 49 | |
| | 40 - 44 | | | | | | | | | | 40 - 44 | |
| | 35 - 39 | | | | | | | | | | 35 - 39 | |
| | 30 - 34 | | | | | | | | | | 30 - 34 | |
| | 25 - 29 | | | | | | | | | | 25 - 29 | |
| | 20 - 24 | | | | | | | | | | 20 - 24 | |
| | Write ≤ 19 | | | | | | | | | | Write ≤ 19 | |
| Respiratory Distress | Severe | | | | | | | | | | Severe | |
| | Moderate | | | | | | | | | | Moderate | |
| | Mild | | | | | | | | | | Mild | |
| | Nil | | | | | | | | | | Nil | |
| O ₂ Saturation (SpO ₂) (%) | ≥ 95 | | | | | | | | | | ≥ 95 | |
| | 92 - 94 | | | | | | | | | | 92 - 94 | |
| | 90 - 91 | | | | | | | | | | 90 - 91 | |
| | Write ≤ 89 | | | | | | | | | | Write ≤ 89 | |
| O ₂ Flow Rate | Write value (L/min) | | | | | | | | | | Write value (L/min) | |
| | Delivery Method | Write | | | | | | | | | Write | |
| Probe Change | Tick | | | | | | | | | | Tick | |
| | Pulse Rate (beats/min) | Write ≥ 190 | | | | | | | | | | Write ≥ 190 |
| 180s | | | | | | | | | | | 180s | |
| 170s | | | | | | | | | | | 170s | |
| 160s | | | | | | | | | | | 160s | |
| 150s | | | | | | | | | | | 150s | |
| 140s | | | | | | | | | | | 140s | |
| 130s | | | | | | | | | | | 130s | |
| 120s | | | | | | | | | | | 120s | |
| 110s | | | | | | | | | | | 110s | |
| 100s | | | | | | | | | | | 100s | |
| Write ≤ 99 | | | | | | | | | | | Write ≤ 99 | |
| Capillary Refill | | Write ≥ 3 sec | | | | | | | | | | Write ≥ 3 sec |
| | | < 3 sec | | | | | | | | | | < 3 sec |
| Blood Pressure (mmHg) | | Write ≥ 110 | | | | | | | | | | Write ≥ 110 |
| | 100s | | | | | | | | | | 100s | |
| | 90s | | | | | | | | | | 90s | |
| | 80s | | | | | | | | | | 80s | |
| | 70s | | | | | | | | | | 70s | |
| | 60s | | | | | | | | | | 60s | |
| | 50s | | | | | | | | | | 50s | |
| | 40s | | | | | | | | | | 40s | |
| | Write ≤ 39 | | | | | | | | | | Write ≤ 39 | |
| | Temp (°C) | Write ≥ 39.1 | | | | | | | | | | Write ≥ 39.1 |
| 38.6 - 39.0 | | | | | | | | | | | 38.6 - 39.0 | |
| 38.0 - 38.5 | | | | | | | | | | | 38.0 - 38.5 | |
| 37.6 - 37.9 | | | | | | | | | | | 37.6 - 37.9 | |
| 37.1 - 37.5 | | | | | | | | | | | 37.1 - 37.5 | |
| 36.6 - 37.0 | | | | | | | | | | | 36.6 - 37.0 | |
| 36.1 - 36.5 | | | | | | | | | | | 36.1 - 36.5 | |
| 35.6 - 36.0 | | | | | | | | | | | 35.6 - 36.0 | |
| Write ≤ 35.5 | | | | | | | | | | | Write ≤ 35.5 | |
| Level of Consciousness (wake patient before scoring) | | Alert | | | | | | | | | | Alert |
| | Verbal | | | | | | | | | | Verbal | |
| | Pain | | | | | | | | | | Pain | |
| | Unresponsive | | | | | | | | | | Unresponsive | |
| Level of Sedation <i>For children receiving sedation and/or opioids only (wake patient before scoring)</i> | 3 | | | | | | | | | | 3 | |
| | 2 | | | | | | | | | | 2 | |
| | 1 | | | | | | | | | | 1 | |
| | 0 | | | | | | | | | | 0 | |
| Pain Score Refer to FLACC score | 8 - 10 | | | | | | | | | | 8 - 10 | |
| | 5 - 7 | | | | | | | | | | 5 - 7 | |
| | 0 - 4 | | | | | | | | | | 0 - 4 | |
| BGL | Write (mmol/L) | | | | | | | | | Write (mmol/L) | | |
| Initials | | | | | | | | | | | | |

| | | | |
|--|--|-------------------|--|
|  <p>Rapid Detection and Response Paediatric Observation Chart (0 - 3 months) MR-59B</p> <p>Hospital/Site:.....</p> | Affix patient identification label in this box | | |
| | U.R. No:..... | | |
| | Surname:..... | | |
| | Given Name:..... | | |
| | Second Given Name:..... | | |
| D.O.B.: | | Sex/Gender: | |

| SECTION D - MODIFICATIONS | | | | |
|--|----------------|----------------|----------------|----------------|
| A Medical Officer must write and review any Modifications . These are any observation(s) for this patient within a specified time that modify the trigger point for escalation. Refer to the local procedure(s) for instructions on documenting and altering Modifications. | | | | |
| | Modification 1 | Modification 2 | Modification 3 | Modification 4 |
| Start Date and Time | | | | |
| Finish Date and Time | | | | |
| Observation(s) | | | | |
| Triggers for MDT review | | | | |
| Triggers for MER call | | | | |
| Doctor's Signature | | | | |
| Doctor's Name (print) | | | | |
| Doctor's Designation | | | | |
| Nurse/Midwife Signature | | | | |
| Nurse/Midwife Name (print) | | | | |
| Nurse/Midwife Designation | | | | |

| SECTION E - FREQUENCY OF OBSERVATIONS | | | | | | | |
|--|----------------------|-----|-----|-----|-----|-----|-----|
| Observations should be performed routinely at least 4 hourly unless advised below. Refer to local procedure for who can alter frequency. | | | | | | | |
| Date | (e.g.) 06/04/2021 | / / | / / | / / | / / | / / | / / |
| Frequency | 2/24 | | | | | | |
| Name/Designation | Smith RN | | | | | | |

| SECTION F - INTERVENTION OR REVIEW DONE (INCLUDING MDT OR MET CALL) | | | | | |
|---|---|-------------------------------|--------------------------|--------------------------|-----------|
| Date | Intervention or review (e.g. Urine Output, increase frequency BGL's, O ₂ changes etc) | Patient family/ carer concern | Physical state change | Mental state change | Name |
| Time | | | | | Signature |
| | | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | |
| | | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | |
| | | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | |
| | | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | |
| | | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | |
| | | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | |