

South Australian Perinatal Practice Guideline

Third and fourth degree tear management

© Department for Health and Wellbeing, Government of South Australia. All rights reserved.

Note:

This guideline provides advice of a general nature. This statewide guideline has been prepared to promote and facilitate standardisation and consistency of practice, using a multidisciplinary approach. The guideline is based on a review of published evidence and expert opinion.

Information in this statewide guideline is current at the time of publication.

SA Health does not accept responsibility for the quality or accuracy of material on websites linked from this site and does not sponsor, approve or endorse materials on such links.

Health practitioners in the South Australian public health sector are expected to review specific details of each patient and professionally assess the applicability of the relevant guideline to that clinical situation.

If for good clinical reasons, a decision is made to depart from the guideline, the responsible clinician must document in the patient's medical record, the decision made, by whom, and detailed reasons for the departure from the guideline.

This statewide guideline does not address all the elements of clinical practice and assumes that the individual clinicians are responsible for discussing care with consumers in an environment that is culturally appropriate and which enables respectful confidential discussion. This includes:

- The use of interpreter services where necessary,
- Advising consumers of their choice and ensuring informed consent is obtained,
- Providing care within scope of practice, meeting all legislative requirements and maintaining standards of professional conduct, and
- Documenting all care in accordance with mandatory and local requirements

Explanation of the aboriginal artwork:

The aboriginal artwork used symbolises the connection to country and the circle shape shows the strong relationships amongst families and the aboriginal culture. The horse shoe shape design shown in front of the generic statement symbolises a woman and those enclosing a smaller horse shoe shape depicts a pregnant women. The smaller horse shoe shape in this instance represents the unborn child. The artwork shown before the specific statements within the document symbolises a footprint and demonstrates the need to move forward together in unison.



Australian Aboriginal Culture is the oldest living culture in the world yet Aboriginal people continue to experience the poorest health outcomes when compared to non-Aboriginal Australians. In South Australia, Aboriginal women are 2-5 times more likely to die in childbirth and their babies are 2-3 times more likely to be of low birth weight. The accumulative effects of stress, low socio economic status, exposure to violence, historical trauma, culturally unsafe and discriminatory health services and health systems are all major contributors to the disparities in Aboriginal maternal and birthing outcomes. Despite these unacceptable statistics the birth of an Aboriginal baby is a celebration of life and an important cultural event bringing family together in celebration, obligation and responsibility. The diversity between Aboriginal cultures, language and practices differ greatly and so it is imperative that perinatal services prepare to respectively manage Aboriginal protocol and provide a culturally positive health care experience for Aboriginal people to ensure the best maternal, neonatal and child health outcomes.

Purpose and Scope of PPG

This guideline provides clinicians with information on the management of third and fourth degree tears. It includes details on risk factors, diagnosis and classification, repair technique and postnatal care and follow-up.



Third and fourth degree tear management

Table I: Complications associated with 3rd and 4th degree tears¹

Serious risks	
> Incontinence of faeces / flatus	Common
> Need for LSCS in future pregnancies due to persistent symptoms of incontinence or abnormal anal sphincter structure or function	Uncommon
> Haematoma	Rare
> Consequences of failure of the repair requiring the need for further interventions e.g. secondary repair or sacral nerve stimulation	Rare
> Rectovaginal fistula	Very rare
Frequent risks	
> Fear, difficulty and discomfort in passing stools in the immediate postpartum period > Migration of suture material requiring removal > Granulation tissue formation	
> Faecal urgency	26/100 Very common
> Perineal pain and dyspareunia	9/100 Common
> Wound infection	8/100 Common
> Urinary infection	

Adapted from RCOG Consent advice No. 9, repair of third and fourth degree perineal tears following childbirth¹⁴

RCOG: Presenting information on risk²

Term	Equivalent numerical ratio
Very common	1/1 to 1/10
Common	1/10 to 1/100
Uncommon	1/100 to 1/1,000
Rare	1/1,000 to 1/10,000
Very rare	Less than 1/10,000

Based on the RCOG Clinical Governance Advice, Presenting information on Risk¹⁵



Third and fourth degree tear management

Table of Contents

[Purpose and Scope of PPG](#)

[Table I: Complications associated with 3rd and 4th degree tears](#)

[Summary of Practice Recommendations](#)

[Abbreviations](#)

[Literature review](#)

[Classification of tears](#)

[Risk factors](#)

[Management of repairs](#)

[Diagnosis of third and fourth degree tear](#)

[Recommended method for repair](#)

[Repair technique](#)

[Recommended antibiotic cover](#)

[Role of colostomy](#)

[Postpartum Management](#)

[Bladder management](#)

[Antibiotics](#)

[Analgesia and other measures](#)

[Follow-up \(at 6 weeks\)](#)

[Recommendations about future pregnancies](#)

[References](#)

[Acknowledgements](#)

Summary of Practice Recommendations

Careful inspection of the perineum, vulva and vagina following birth is essential

Rectal examination prior to suturing is required when the woman has undergone episiotomy or if the tear extends to the anal verge to determine classification of tear

Repair of 3b, 3c and fourth degree tears should be undertaken in theatre with adequate analgesia

Consult a colorectal surgeon if a large fourth degree tear is diagnosed

Antibiotic cover is required for all third and fourth degree tears

Postnatal bladder management requires specific attention

Postnatal follow-up needs to be individualised

Recommendations for subsequent births is based on presence of symptoms and the woman's preference

Abbreviations

et al.	And others
EAS	External anal sphincter
e.g.	For example
g	Gram(s)
IAS	Internal anal sphincter
IV	Intravenous
mg	Milligrams
n	Number
%	Percent
®	Registered trademark
RCOG	Royal College of Obstetricians and Gynaecologists



Third and fourth degree tear management

Literature review

- > Obstetric anal sphincter injury encompasses both third and fourth degree perineal tears and can occur with an intact perineum
- > In South Australia in 2010, third and fourth degree tears occurred in 3.4 % (n=450) of vaginal births³
- > Prospective studies using postpartum anal endoanal sonography suggest that almost one third of primiparous women may sustain occult anal sphincter injury following vaginal birth^{4,5,6}
- > Recent randomised controlled studies of external anal sphincter (EAS) repair have reported low incidences of anal incontinence symptoms (e.g. loss of control over flatus, faecal urgency and staining) with 60 – 80 % of women asymptomatic at 12 months⁵
- > Damage to the innervation of the sphincter muscles and pelvic floor may be related to pudendal nerve damage^{4,7}

Classification of tears

First Degree:

- > Injury to the perineal skin only

Second Degree:

- > Injury to the perineum extending into the perineal muscles but not the anal sphincter (either external [EAS] or internal anal sphincter[IAS])

Third degree:

- > Injury to the perineum involving the anal sphincter complex:
 - > 3a: Less than 50 % of EAS thickness torn
 - > 3b: More than 50 % of EAS thickness torn
 - > 3c: Both EAS and IAS torn⁴

Fourth Degree:

- > Disruption of the anal sphincter complex (EAS and IAS) and anal epithelium⁵. Occasionally there can be an anal or rectal mucosa tear behind an intact sphincter. Rectal examination before repair is recommended⁸

Risk factors

- > First vaginal birth
- > Instrumental delivery
- > Prolonged second stage
- > Macrosomia > 4 kg
- > Midline episiotomy
- > Occipitoposterior position at delivery
- > Induction of labour
- > Epidural analgesia
- > Shoulder dystocia
- > Most of the above risk factors cannot readily be used to prevent or predict the occurrence of a third or fourth degree tear⁹
- > Damage to the pudendal nerves is cumulative in successive vaginal births^{7,10,11}



Third and fourth degree tear management

Management of Repair

- > All women should be examined following vaginal birth to assess the degree of vaginal, perineal or rectal injury
- > All sphincter damage must be identified, documented and treated appropriately
- > This includes:
 - > partial sphincter tears
 - > sphincter damage with an intact perineum
 - > "buttonhole" rectal mucosa tears
- > Accurate diagnosis will mean that these women will have the best chance of normal anal function in years to come

Diagnosis of third and fourth degree tear

All women delivering vaginally should have:

- > Informed verbal consent explaining the need for thorough examination of the vagina, vulva and perineum and why a per rectum examination may be required
- > Good exposure and good lighting
- > Good analgesia
- > Vulval and vaginal examination
- > The normal pattern of peri-anal rugae confirmed
- > Rectal examination for all episiotomies or if tear extending to anal verge
- > Direct visualisation of sphincter with digit in rectum
- > Palpation of sphincter with digit in rectum and pill rolling action with thumb on sphincter

Recommended method for repair

- > Third and fourth degree repairs should be undertaken by an obstetrician or a registrar trained to repair third and fourth degree tears after discussion with a consultant
- > 3a tears may be repaired in labour and delivery if there is adequate analgesia
- > All 3 b and c and fourth degree tear repairs should be carried out in theatre with adequate regional anaesthesia to facilitate adequate analgesia, good visualisation and relaxation of sphincter muscles

Repair technique

- > Perform a repeat detailed assessment of the degree of vaginal / perineal / rectal injury under anaesthesia
- > Ends of EAS should be mobilised by sharp and blunt dissection to facilitate a tension free repair
- > When repairing the EAS, use either monofilament sutures such as 2-0 polydioxanone (PDS or Maxon) (DON'T USE Vicryl)
- > For repair of complete tear of the EAS, either an overlapping or end-to-end (approximation) method can be used. Overlapping repair is preferred by most Obstetric Consultants specialising in the management of anal sphincter injury and Colorectal Surgeons; however, there is no level I evidence to support this⁶
- > Where the IAS can be identified, it is advisable to repair separately with interrupted sutures. When repairing the IAS, use fine suture size such as 3-0 PDS[®] and 2-0 Vicryl[®] (associated with less irritation and discomfort)
- > 3a tears can only be repaired using an end-to-end repair
- > Bury surgical knots beneath the superficial perineal muscles by performing a standard perineal repair to prevent knot migration to the skin
- > Perform a rectal examination on completion to ensure the repair is intact
- > Document the procedure in case notes and arrange postpartum follow up



Third and fourth degree tear management

Recommended antibiotic cover

- > Give single IV doses of both cephazolin 2 g and metronidazole 500 mg

Allergy to penicillin

- > Single IV doses of clindamycin 450 mg, AND gentamicin 5 mg / kg

Role of colostomy

- > There is no clear consensus amongst colorectal surgeons on who requires Colostomy and no reliable data to base a decision on
- > A Colostomy is not required for management of 3a, 3b and straightforward 3c tears
- > A Colostomy is usually indicated with large 4th degree tears, especially when the tear extends above the levator muscles, or where other risk factors for fistula exist
- > Consult with a colorectal surgeon regarding the need for a colostomy

Postpartum Management

Bladder management

- > On average, bladder sensation takes between 6 to 7 hours to return after a vaginal birth with regional anaesthesia¹²
- > In cases of 3rd or 4th degree tear, severe perineal discomfort is known to cause urinary retention with a delay of up to 12 hours before bladder sensation returns¹³
- > Urinary catheterisation should occur following 3rd and 4th degree repair in the immediate postpartum period to minimise urinary retention. The optimum time for catheterisation after birth is uncertain. Careful attention should be paid to voiding after removal of the catheter, particularly in the first six hours after catheter removal (see *Bladder Management for Intrapartum and Postnatal Women* PPG available at www.sahealth.sa.gov.au/perinatal)

Antibiotics

- > The use of broad-spectrum antibiotic cover is recommended after obstetric anal sphincter repair to reduce the incidence of postoperative infections and wound dehiscence⁶
- > Commence oral Augmentin Duo Forte® (amoxicillin 875 mg and clavulanic acid 125 mg) 12 hourly with meals for 5 days
- > If allergic to penicillin, use both
 - > oral ciprofloxacin 500 mg 12 hourly for 5 days
 - plus
 - > oral clindamycin 450 mg 8 hourly for 5 days

Breastfeeding: All these drugs are acceptable

Analgesia and other measures

- > Use a multimodal approach to minimise the use of opioid medication, i.e. oxycodone and codeine containing analgesics, as they may cause constipation
- > Administer oral paracetamol 1 g every 6 hours as required
- > If there are no contra-indications, administer diclofenac (Voltaren®) 100 mg suppository per rectum at the end of the procedure while the patient is in the lithotomy position. Subsequent doses can be administered orally (i.e. 50 mg TDS), commencing no sooner than ten hours after administration of the intra-operative dose
- > Ice packs and resting supine / prone for 10 - 20 minutes every 2 – 3 hours over the first week may decrease symptoms of pelvic floor fatigue (e.g. swelling, pain and perineal descent)
- > Bulking agents and stool softeners (e.g. Fybogel® 1 sachet three times a day, Lactulose® 20 mL twice daily, and Coloxyl® 120 mg 1-2 nocte in addition as required) are recommended. Commence after 24 hours and continue for two weeks before weaning off. Educate the woman about the need for adequate fluid intake when using bulking agents
- > In-patient referral to a continence health professional (e.g. continence nurse advisor /

Third and fourth degree tear management

practitioner or physiotherapist) for advice about defecation techniques, pelvic floor care and ongoing support

- > Before discharge, the woman must be fully informed about the nature of her injury, associated risks (see Table 1) and benefits of follow-up

Follow – up (at 6 weeks)

- > A third or fourth degree tear is a significant peripartum event. Postpartum follow-up by a consultant with an interest in management of third and fourth degree tears and a continence health professional referral is required. If the woman is experiencing incontinence or pain at follow-up, a colorectal opinion and investigation (endoanal ultrasound) may be necessary
- > Establish the following:
 - > Control of bowel motions
 - > Control of flatus
 - > Faecal urgency
 - > Offensive vaginal discharge (this may suggest a fistula)
 - > Confirm urinary continence
 - > Assess pelvic floor muscles
 - > Assess ongoing perineal discomfort
- > The mode of subsequent delivery should be discussed in the context of current symptoms or findings of postpartum sonography

Recommendations about future pregnancies

- > Women who are asymptomatic may consider a vaginal birth
- > Advise the woman that there is no evidence to support the role of prophylactic episiotomy in subsequent pregnancies
- > Recommend LSCS:
 - > Symptomatic
 - > Previous 4th degree tear
 - > Delayed surgical correction of sphincter damage
 - > Other risk factor for sphincter damage (e.g. big baby, occipito posterior position)
 - > Woman's request
- > It is appropriate to warn women that the cumulative effect of ageing, menopause and progression of neuropathy on long term sphincter weakness by the fifth and sixth decade may result in the new onset of symptoms for which treatment is available⁵



Third and fourth degree tear management

References

1. Royal College of Obstetricians and Gynaecologists. Repair of third and fourth degree perineal tears following childbirth. Consent advice No 9. June 2010. Available from URL: <http://www.rcog.org.uk/repair-third-and-fourth-degree-perineal-tears-following-childbirth>
2. Royal College of Obstetricians and Gynaecologists. Presenting information on risk. Clinical Governance advice No. 7. London: RCOG; 2008. Available from URL: <http://www.rcog.org.uk/files/rcog-corp/CGA7-15072010.pdf>
3. Scheil W, Scott J, Catcheside B & Sage L. Pregnancy Outcome in South Australia 2010. Adelaide: Pregnancy Outcome Unit, SA Health, Government of South Australia, 2012.
4. Sultan AH, Kamm MA, Hudson CN, Chir M, Thomas JM, Bartram CI. Anal-Sphincter disruption during vaginal delivery. N Engl J Med 1993; 329: 1905-11 (Level III).
5. Royal College of Obstetricians and Gynaecologists (RCOG). Management of third and fourth-degree perineal tears following vaginal delivery, Guideline No. 29. London: RCOG Press; 2007.
6. Thakar R, Sultan AH. Management of obstetric anal sphincter injury. The Obstetrician and Gynaecologist 2003; 5: 72-8.
7. Fitzpatrick M, O'Herlihy C. The effects of labour and delivery on the pelvic floor. Best Practice & Research Clinical Obstet Gynaecol 2001; 15: 63 – 79.
8. Signorello LB, Harlow BL, Chekos AK, Repke JT. Midline episiotomy and anal incontinence: Retrospective cohort study. Br Med J 2000; 320: 86 – 91 (Level III-2).
9. Rieger N, Perera S, Stephens J, Coates D, Po D. Anal sphincter function and integrity after primary repair of third-degree tear: a prospective analysis. ANZ J of Surg 2004; 74 122 – 4 (Level III-3).
10. Fynes M, Donnelly V, Behan M, O'Connell PR, O'Herlihy C. Effect of second vaginal delivery on anorectal physiology and faecal continence: a prospective study. The Lancet 1999; 354:983-86 (Level III).
11. Fynes M. Childbirth and faecal incontinence. Aust Continence Journal 2001; 7: 2-6 (Level III).
12. Foona R, Toozs-Hobsonb P, Millnsb P, Kilbyb M. The impact of anesthesia and mode of delivery on the urinary bladder in the postdelivery period. Int J Gyn Obst 2010; 110: 114-117.
13. Sultan AH, Thakar R, Fenner DE, editors. Perineal and anal sphincter trauma. London: Springer; 2009.

Useful resources

- > Royal College of Obstetrics and Gynaecology Patient information Third and fourth degree tears. Available from URL: <https://www.rcog.org.uk/en/patients/patient-leaflets/third--or-fourth-degree-tear-during-childbirth/>
- > Royal Women's Hospital. Consumer information. Anal sphincter tears in childbirth. Available from URL: <https://thewomens.r.worldssl.net/images/uploads/fact-sheets/Perineal-tears-third-and-fourth-degree.pdf>



Third and fourth degree tear management

Acknowledgements

The South Australian Perinatal Practice Guidelines gratefully acknowledge the contribution of clinicians and other stakeholders who participated throughout the guideline development process particularly:

Write Group Lead

Dr Steven Scroggs

Write Group Members

Allison Rogers
Dr Brian Peat
Dr Elinor Atkinson
Prof Gus Dekker

Other major contributors

J Phelps
J Turnidge
N Rieger
SAPPG Work Group 2004-2014

SAPPG Management Group Members

Sonia Angus
Dr Kris Bascomb
Lyn Bastian
Elizabeth Bennett
Dr Feisal Chenia
John Coombas
A/Prof Rosalie Grivell
Dr Sue Kennedy-Andrews
Jackie Kitschke
Catherine Leggett
Dr Anupam Parange
Dr Andrew McPhee
Rebecca Smith
A/Prof John Svigos
Dr Laura Willington



Third and fourth degree tear management

Document Ownership & History

Developed by: SA Maternal, Neonatal & Gynaecology Community of Practice
Contact: HealthCYWHSPerinatalProtocol@sa.gov.au
Endorsed by: SA Safety and Quality Strategic Governance Committee
Next review due: 17 June 2019
ISBN number: 978-1-74243-162-8
PDS reference: CG143
Policy history: Is this a new policy (V1)? **N**
Does this policy amend or update an existing policy? **Y**
If so, which version? **V5**
Does this policy replace another policy with a different title? **N**
If so, which policy (title)?

Approval Date	Version	Who approved New/Revised Version	Reason for Change
15 June 2018	V5.1	SA Health Safety and Quality Strategic Governance Committee	Review date extended to 5 years following risk assessment. New template.
17 June 2014	V5	SA Health Safety and Quality Strategic Governance Committee	Reviewed in line with scheduled review date
20 Mar 2012	V4	Maternal and Neonatal Clinical Network	Reviewed in line with scheduled review date
27 Oct 2009	V3	Maternal and Neonatal Clinical Network	Reviewed in line with scheduled review date
25 Jan 2005	V2	Maternal and Neonatal Clinical Network	Reviewed.
21 Jul 2004	V1	Maternal and Neonatal Clinical Network	Original approved version.

