Vascular

Outpatient service information, triage and referral guidelines

Description of Service:

Vascular services for Central and Northern Adelaide is based at the Royal Adelaide Hospital, but consults and operates at several sites across the networks based on disease complexity and patient needs.

Our outpatient services are multi-disciplinary units which provide assessment, management and treatment for all subspecialty areas of the vascular system.

Services Provided:		Exclusions:	
>	Symptomatic varicose veins	>	Paediatrics
>	Venous and arterial ulcers	>	Cosmetic varicose veins
>	Severe limb ischaemia and claudication		
>	Abdominal aortic aneurysms		
>	Complicated and diabetic foot conditions		
>	Vascular renal access		

For admission or URGENT advice contact the Vascular Registrar									
Royal Adelaide Hospital (RAH)	Ph: (08) 7074 0000								
Alternatively, if the problem is urgent and the patient is unstable, the patient should be sent directly to the									
Emergency Department of the nearest hospital for assessment.									
For more information or to make a referral contact									
Royal Adelaide Hospital (RAH)	Fax: (08) 7074 6247 Ph: 1300 153 853								
Essential Referral Content:									
Demographic	Clinical								
> Date of birth	> Reason for referral								
> Contact details (including mobile phone)	> Duration of symptoms								
> Referring GP details	 Relevant pathology and imaging reports 								
> Interpreter requirements	> Past medical history								
> Medicare number	> Current medications								



Adult Triage Criteria for referral									
Emergency Referrer must contact Vascular on-call registrar at the RAH to discuss care or arrange potential urgent review via an emergency department	Category 1 Up to 30 days	Category 2 Up to 90 days	Category 3 Greater than 90 days						
 > AAA with history of back pain, abdominal pain or tenderness > AAA >7cm + symptoms > TIA/Stroke symptoms within 2 weeks > Acutely Ischaemic foot > Diabetic foot infection with erythema >2cm from ulcer > Iliofemoral/Upper DVT – with confirmed duplex scan > Carotid dissection > Aortic dissection > Superficial thrombophlebitis extending to within 5cm of the SFJ or SPJ and confirmed on ultrasound scan 	 Neuropathic ulceration Infected venous leg ulceration Carotid body tumour Asymptomatic AAA >5cm Ischaemic rest pain Foot ulceration Require appointment within 1 week: TIA/stroke within 2 week symptom period (combined with referral to TIA clinic)	 > Arterio-venous malformation. > AAA <5cm > Vasospastic disorders > Short distance claudication <50cm > Chronic venous leg ulceration 	 Hyperhidrosis Claudication Thoracic outlet syndrome – arterial, venous, neurogenic Distal superficial thrombophlebitis (confirmed with duplex scan) Varicose veins Asymptomatic carotid disease. Note: varicose vein inclusion criteria Current or healed ulcers Skin changes in keeping with chronic venous insufficiency Bleeding varicosity Phlebitis or thrombosis confirmed with ultrasound.						

For more information or to make a referral

Royal Adelaide Hospital: Level 3F, Wing 1 Referral Fax Number: (08) 7074 6247 Phone Number: 1300 153 853 Acknowledgement: This document has had the final approval from the head of unit: Vascular Department

Version	Approved date	Review date	Amendment
Draft V0.1	20192502	20202502	

Web: www.sahealth.gov.au

 $\ensuremath{\textcircled{\sc be}}$ Department for Health and Wellbeing, Government of South Australia. All rights reserved. 18020

