



Your guide to pregnancy

Flinders Women and Children



Government
of South Australia

Health

Southern Adelaide
Local Health Network

List of important telephone numbers you may need

Flinders Medical Centre (main switchboard-24 hours)	8204 5511
Admission Office	8204 4126
Women's Health Clinic (between 10am - 4pm)	8204 5197
Women's Health Clinic midwife	8204 4667 8204 4680
Birthing and Assessment Suite (phone switchboard)	8204 5511
Birth Centre	8204 3130
Physiotherapy Department	8204 5498
Antenatal Ward	8204 4217
Postnatal Ward	8204 4216
Maternity Outreach	8204 5189
Postnatal Support Service	8204 4216
Neonatal Unit	8204 5041
Paediatric Unit	8204 4629
Noarlunga Health Services	8384 9222
Noarlunga Maternity Care Clinics	8384 9454

Welcome to Flinders Medical Centre

Congratulations and best wishes for the coming birth of your baby.

Pregnancy and childbirth are very special events. They bring with them joy and new responsibilities for a woman and her family, but may also present with uncertainty and anxiety. The aim of this booklet is to reassure women, by providing information about pregnancy, childbirth and their stay in Flinders Medical Centre.

It is a philosophy of Flinders Medical Centre, Women's and Children's Division that midwives provide and maintain a relaxed environment and an atmosphere in which women feel supported physically, emotionally and spiritually throughout their pregnancy and birth journey.

Obstetric antenatal care is provided by doctors and midwives at Flinders Medical Centre and at Noarlunga Health Service, Maternity Care clinics.

A comprehensive range of supporting services are available to ensure pregnancy is an enjoyable time and the birth of your baby a safe and memorable event.

Flinders Medical Centre is a Baby Friendly Health Initiative accredited hospital and, as such, supports and promotes breast feeding.



Contents

Antenatal Information	6	General care in pregnancy	9
Antenatal clinic appointment	6	Dental	9
Importance of antenatal care	6	Medication	9
Pregnancy records	6	Drugs and Alcohol	9
Models of Antenatal Care	6	Smoking	9
Obstetric antenatal care with a doctor	6	Rest	9
Midwives clinics	6	Nutrition	9
Southern Midwifery Group Practice	6	Foods to avoid	10
Shared Care	7	Contact with illness	10
Karpa Ngawattendi (Aboriginal Health Unit)	7	Sexual intercourse	10
Antenatal testing	7	When to contact the hospital	10
Blood & Urine test	7	Common discomforts	11
First Trimester Screening	7	Nausea/vomiting	11
Second Trimester Screening	7	Constipation	11
NIPT (Non-Invasive Antenatal Testing)	8	Haemorrhoids	11
Routine Investigations in Pregnancy	8	Varicose veins	11
Other Health Professionals involved in Antenatal Care	8	Backache	11
Fitness during pregnancy	9	Reflux (heartburn)	11
		Cramps	11
		General Information	12
		Admission to Flinders Medical Centre	12
		What to bring in to Hospital:	12
		What to bring for Baby	12

Home Preparation	12	Infant feeding	17
Practical tips	12	Breast Feeding	17
Baby safe capsule	13	Artificial feeding	17
Labour	13	Registration Forms	17
Early labour at home	13	Neonatal (Baby) examinations and tests	18
When labour begins – what to do	14	Community Support	18
Breaking of Membranes (waters)	14	Maternity Outreach	18
Going to Hospital - Birthing and Assessment Suites	14	Post-natal day service	18
Birthing and assessment Suite	14	Child and Youth Health Services (CYH)	18
Support during labour	14	Australian Breast Feeding association (ABA)	18
Care during labour	15	Multiple Birth Association	18
Pain relief	15	Postnatal Depression	19
Induction of Labour	15	Resources	20
Assisted delivery	15	Websites	20
Episiotomy	15		
Forceps and Ventouse delivery	15		
Caesarean section	15		
After the birth of the baby	16		
Postnatal Ward and Care	16		
Visiting hours	16		
Neonatal Unit	17		
Discharge from Hospital	17		
Physiotherapy	17		

Antenatal Information

Antenatal clinic appointment

Antenatal clinic appointments are attended at either Flinders Medical Centre in Women's Health Clinic level 1 (ground floor) or at Noarlunga Health Service in the Maternity Care clinic, outpatients.

Importance of antenatal care

Pregnancy and Childbirth are promoted as a normal process. Regular antenatal checks throughout the pregnancy are required to ensure you and your baby are healthy. One person is welcome to attend the antenatal appointments with you, either partner or support person.

Covid-19 has created restrictions in persons coming to antenatal clinics both at Flinders Medical Centre and Noarlunga Health Service. Please phone Women's Health Clinic 8204 4667 to obtain the latest visitor outpatient restrictions.

Pregnancy records

You will receive an orange folder; South Australian Pregnancy Record at your first visit, if not already given by the GP (local doctor). It is important the South Australian Pregnancy Record is bought to each appointment as it contains information regarding your pregnancy recorded at each visit with all clinicians.

Models of Antenatal Care

During the first obstetric appointment (Pre-Obstetric booking - POB), models of antenatal care will be discussed to consider depending on individual medical and obstetric histories. Types of antenatal care include a medical model with a doctor, low risk antenatal care with a midwife, shared care with a registered shared care GP (general practitioner) or southern midwifery group practice, which is midwifery based only with strict low risk criteria.

Obstetric antenatal care with a doctor

This model of care usually is for women who may have some prior medical or obstetric history that requires specialist care.

Midwives clinics

Midwives clinics are designed for low risk women who would like the options of all types of pain relief in labour or have an obstetric history that prevents them from being in Southern Midwifery Group Practice such a previous caesarean section.

Southern Midwifery Group Practice

Southern Midwifery Group Practice is an antenatal model of care designed at low risk women who do not have any medical or complicating obstetric history. Ideally these women would like to deliver their baby with minimal or no pain relief and be discharged home within 24 hours of the birth.

Shared Care

Shared care is antenatal care shared with your local GP who must be accredited with the South Australian Shared Care programme. The majority of the antenatal care will be with the GP. Towards the end of the pregnancy an antenatal appointment is required at Flinders Medical Centre for continuity of care within the hospital system.

Karpa Ngawattendi (Aboriginal Health Unit)

This is a free service for Aboriginal women living in Southern Metropolitan Adelaide who plan to birth at Flinders Medical Centre.

Phone: 8204 6359

Antenatal testing

Blood & Urine test

Usually when pregnancies are diagnosed, women attend a GP appointment and initial testing is started.

This usually comprises of a blood test, testing for a variety of blood levels including Blood Group and Antibodies, Full Blood Count, Hepatitis B&C, HIV and Syphilis infections and a Midstream Urine specimen.

The benefit of these tests are that conditions can be treated which can improve prognosis and prevent transmission of a virus/infection to the baby, should a positive result occur.

Women who have Rhesus negative blood groups are recommended to have prophylactic Anti D at 28 and 34 weeks gestation, after birth and after any bleeding occurrence during pregnancy. This is to prevent positive antibodies being made in the mother's blood.

First Trimester Screening

First Trimester Screening is optional but is recommended and requires to be ordered by your GP. This test comprises of a blood test and ultrasound performed between 11+0 and 14+0 weeks (the blood test can be performed as early as 9+0 weeks). This test ascertains the screening risk of certain chromosomal abnormalities for your baby. These tests are only screen tests and are not diagnostic. Should the test return a high risk reading for chromosomal abnormalities an appointment will be made with the Antenatal Diagnostic Clinic to discuss options.

Second Trimester Screening

This is a blood test only between 14+1 weeks and 20+6 weeks. This test screens for chromosomal abnormalities and Neural Tube Defects and is also an optional test. This screening test is not as effective as the First Trimester Screening, but maybe an option if pregnancy is discovered late. Should this screening test return a positive high risk result for chromosomal abnormalities, an appointment will be made with the Antenatal Diagnostic Clinic to discuss options.

NIPT (Non-Invasive Antenatal Testing)

The GP may offer a NIPT test to determine the risk of chromosomal abnormalities. This is a blood test only. This test is quite specific for certain chromosomal abnormalities, however is a screen test only. This is quite an expensive test, with no Medicare rebate. Please discuss with your local doctor or request to speak with a clinician at Flinders Medical Centre for further information. Should the test return a high-risk result for chromosomal abnormalities an appointment will be made with the Antenatal Diagnostic Clinic to discuss options.

Routine Investigations in Pregnancy

Morphology scan

Usually performed between 19 – 22 weeks gestation. The ultrasound assesses the baby's growth, structural development and detects the placenta and cord location.

Oral Glucose Tolerance Test (OGTT)

This test is performed at 28 weeks gestation and assesses for Gestational Diabetes. Gestational diabetes is a common condition that can occur during pregnancy and usually resolves after birth of the baby and placenta. If you have risk factors for gestational diabetes, an early OGTT will be arranged around 16 weeks gestation.

Other routine tests at 28 weeks include Haemoglobin (iron) levels, blood antibodies and if required, vitamin D levels.

Group B Streptococcus swab (GBS)

This swab is taken at 36 weeks gestation. Should GBS be present, intravenous antibiotics are arranged for labour or if membranes rupture early. Further information will be discussed at your antenatal visits.

Other Health Professionals involved in Antenatal Care

Other health professional available to pregnant women at Flinders Medical Centre include Social Workers, Physiotherapists, Interpreters, Perinatal Mental Health Nurse, Drug and Alcohol support (DASSA) and Aboriginal Health workers.

Flinders Medical Centre is a teaching hospital. Medical and Midwifery students may be present during your care. Any concerns, please discuss with your midwife or doctor.

Childbirth education is on-line, offering tour videos and references to antenatal information and education. There are structured question and answer "Q&A" sessions available. Please visit the Pregnancy Antenatal education pages on the Flinders Medical Centre website prior to booking into a session. Book into a session if you still have questions regarding your birth at Flinders Medical Centre, by calling the Women's Health Clinic on 08204 4680, otherwise refer to the education pages at the back of the orange South Australian Pregnancy Record.

Fitness during pregnancy

Keeping fit during pregnancy will assist in coping with the physical and emotional strain on your body. Back and pelvic joints loosen during pregnancy. Physiotherapists can be referred to aid in any discomfort. The physiotherapists may also assist in other common issues such as abdominal muscle weakness, pelvic floor weakness and bladder control problems.

The Physiotherapy Department provides pregnancy and postnatal exercise classes. Please refer to the information leaflet in the orange South Australia Pregnancy Record.

General care in pregnancy

It is important to attend all antenatal appointments to ensure wellbeing of you and your baby. A normal lifestyle can be maintained in pregnancy, but issues need to be considered.

Dental

Dental appointments can still be attended to in pregnancy.

Medication

Prescribed medication and non-prescribed medication, including herbal medicines or remedies, needs to be discussed with the doctor, midwife or GP, as they may cause harm in pregnancy.

Drugs and Alcohol

Illicit drugs and alcohol are extremely harmful in pregnancy and should be avoided. If you have been taking illicit drugs or alcohol in your pregnancy, please discuss

this with your doctor, midwife or GP and request drug and alcohol support (DASSA).

Smoking

Smoking is a health hazard and women are advised to not smoke during pregnancy. If assistance is required to quit or reduce the smoking number, please discuss with your doctor, midwife or GP for assistance and the smoking Quitline.

Rest

Rest is important, especially later in your pregnancy to lessen fatigue. Try lying on your side for a half an hour each day or try and try to go to bed earlier.

Nutrition

Good food during pregnancy is important in supplying energy and nutrition to yourself and your baby. This does not mean to eat more, but to eat to choose a well balance diet of the 5 main food groups:

- Breads, cereals, pasta, rice (preferably brown for all)
- Fruits and vegetables
- Meat, legumes, fish, eggs, nuts and tofu
- Milk, yoghurt, cheese
- Fats and oils

Extra iron is usually required during pregnancy. The best dietary sources of iron are lean red meat, chicken and fish. Other sources include fortified cereals, green leafy vegetables, eggs, legumes, baked beans, soya beans, almonds, nuts mixed dried fruit and peanut butter. Foods high in vitamin C can aid in absorption of iron, for example orange juice and tomatoes.

Calcium is important for healthy teeth and bone growth and can be obtained in dairy foods. (Please Note: Calcium can interfere with iron absorption so avoid calcium rich food when taking iron supplements).

An iodine supplement (150 micrograms daily) is recommended during pregnancy and whilst breast feeding to enhance growth and development of baby's brain and other organs.

Note: for safety reasons please do not take any vitamin or supplement unless it is discussed with a health professional first.

Foods to avoid

Foods that have the potential to be contaminated with *Listeria* need to be avoided in pregnancy. *Listeria* is a bacterial infection caused by eating certain foods such as deli meats or unpasteurised dairy foods. In pregnancy *Listeria* infection is likely to cause only mild symptoms to the mother but the consequences on the baby can be devastating.

Listeria is destroyed by heat, but can survive in the refrigerator (unlike most bacteria). Eating freshly prepared, freshly cooked foods is the best way to avoid contact with *Listeria*.

Foods to avoid during pregnancy are mostly chilled, ready to eat foods, deli meats and unpasteurised dairy foods. These include:

- Soft cheeses (brie, camembert, ricotta and fetta)
- Takeaway cooked chicken
- Cold meats
- Pate
- Prepared and stored salads
- Raw seafood (oysters, sashimi)
- Smoked seafood (smoked salmon, smoked oysters)
- Soft serve ice cream

For more information about the effects of *Listeria* in pregnancy, ask your doctor or midwife.

Contact with illness

Avoid close contact with people or animals suffering from any kind of illness or infection. Should contact occur, please advise your doctor, midwife or GP.

Sexual intercourse

Sexual intercourse may continue during your pregnancy unless you have been advised otherwise.

When to contact the hospital

Contact Flinders Medical Centre should you experience:

- Bleeding from the vagina
- Vomiting and/or abdominal pain
- Regular contractions before 37 weeks gestation
- Severe headache or visual disturbances (dimness or flashes of light)
- Urinary problems of any kind
- A change in your baby's movements
- If your membranes (waters) break
- If you are concerned or anxious

Phone: (08) 8204 5511 - ask for Birthing and Assessment Suites (BAS) for further advice.

Common discomforts

Nausea/vomiting

This usually occurs during the first three months but can occur for longer in pregnancy.

Try to eat small frequent meals and stay hydrated. Crackers, toast and sandwiches may be best tolerated. Avoid fatty foods or smells that enhance nausea.

Re-hydration may be required in hospital should dehydration become severe.

Constipation

Constipation can be common in pregnancy. Consume a diet rich in fibre, such as wholegrain bread and cereals. Eat plenty of fresh fruit and vegetables or dried fruit. Exercise and drink plenty of water, 6-8 glasses per day and avoid more than 3-4 cups of coffee. Avoid laxatives unless prescribed by your doctor.

Haemorrhoids

Haemorrhoids may appear or become worse if already present before pregnancy. It is important to avoid constipation and straining. Ensure a high fibre diet and drink plenty of water. If this becomes a problem, please discuss with your doctor or midwife.

Varicose veins

Legs and vulval area

Varicose veins may appear or become worse during pregnancy in legs and vulval area. Lying down two to three times a

day, support stockings and ice packs may assist in discomfort and reducing swelling. If these are not effective, a referral to the physiotherapist can be arranged.

Backache

Backache is common in pregnancy. This is due to increase softening of joints and the increasing weight of the growing baby altering posture. This makes women more vulnerable to strain during daily activities. Maintaining a good posture, practicing regular antenatal exercises and wearing comfortable flat-heeled shoes can help to relieve symptoms. If backache persists, please speak to a midwife to organise a tubi-grip support band or a referral to the physiotherapist.

Reflux (heartburn)

Heartburn may develop during pregnancy due to internal organs being moved to make room for the baby. Particular foods such as, acidic, spicy, fatty or rich foods may aggravate the symptoms and should be avoided. Sleep with an extra pillow under your head and shoulders and avoid lying flat. Should reflux persist, speak to you doctor or midwife to recommend reflux relieving medication.

Cramps

Muscle cramps, usually in the legs at night are common in pregnancy. Leg cramps may be relieved by rubbing the affected area or by applying warmth. Calf stretches prior to going to bed may be helpful. If leg cramps continue please discuss with the doctor or midwife.

General Information

Admission to Flinders Medical Centre

All women who are to be admitted to Birthing and Assessment Suite or Southern Midwifery Group Practice are requested to enter via the Emergency Department.

What to bring in to Hospital:

- Supports you might like to use during labour e.g. music
- Orange South Australian Pregnancy Record
- Comfortable casual clothes (shirts that open in the front for ease when breast feeding)
- Supportive maternity bras
- Nightwear
- Dressing gown and slippers
- Toiletries (shampoo/conditioner, soap, toothpaste/toothbrush, brush)
- 1 x box of tissues
- 3-4 packets of sanitary pads
- 1 x box of breast pads
- Underpants
- Pen
- Own pillow if you wish

If choosing to bottle feed or artificially feed (formula), bottles and formula will be provided during your hospital stay, but also bring a tin of your chosen formula for demonstration purposes.

We advise no valuables to be bought into hospital, such as jewellery or large amounts of money.

For safety reasons please do not bring in electrical equipment. TV, radio facilities and hairdryers are available.

What to bring for Baby

- 2 x singlets
- 2 x grow suits or outfits
- 1 x baby jacket
- 1 x blanket
- 2 x wraps
- Mittens
- 1-2 large packets of disposable nappies (newborn size) or fasteners/nappy pins for cloth nappies, including pilchers x 6 (not plastic).
- Baby capsule or car seat for baby aged 0-4 for transport home.

Home Preparation

A new baby can be tiring and emotional for both parents. Caring for a newborn baby is very time consuming. Being prepared and planning for home can help the transition of a new baby easier.

If possible, arrange to have some help with the housework for the first few weeks. Preparing meals which can be frozen and accepting offers of help from family members is advised.

Practical tips

- Arrange the fitting of the baby capsule or car seat to your car, a few weeks prior to the due date
- Prepare some meals in advanced and freeze

- Stock the pantry
- Prepare the baby room and wash all the baby's clothes a few weeks prior to the due date
- Encourage your partner's to perform domestic duties
- Be aware that you may not be able to keep up previous standards of tidiness in the home

Talk to your partner about both your feelings and how you both may be able to assist each other in the adjustment of a new baby.

Take time out for each other. If you have other children, arrange well in advanced of your due date, care arrangements for when you come into hospital.

Baby safe capsule

Legally it is a requirement to have a baby appropriately restrained whilst travelling in a motor vehicle in a baby capsules or 0-4 car seat. 0-4 car seats are available for purchase from department stores or baby furniture stores. Alternatively, baby capsules can be hired.

Labour

Labour is a process that involves three stages. First stage involves the contraction of the uterus and dilation of the cervix. The second stage is when your cervix is fully dilated, and birth of the baby and third stage is the passing of the placenta.

Early labour at home

When contractions start, they might be irregular in strength and length. If possible, stay active by going for a walk or continuing with daily activities. Watching TV or reading a book may distract from the early process. If contractions start at night, it's important to keep relaxing and try and get some sleep, as this process may go on for a while.

Contractions will increase in strength and length and become regular. It is good to have your partner or support person close or with you, in case you require assistance. There are many ways to help assist with the discomfort at home. Try:

- Warm water in the shower or bath (ensure your partner or support person is with you, in case you feel faint or require help getting out of a bath)
- Hot water bottle or heat pack may provide some relief in the lower back or abdomen area.
- Massage to the lower back (or shoulders for distraction)
- Leaning forward into a bean bag or chair to relieve pressure off your back
- Different positions such as lying down, walking or squatting
- Simple analgesia such as Panadol may give some relief (aspirin or ibuprofen cannot be taken)

When labour begins – what to do

This is when contractions are painful, regular, lasting 1-2 minutes. Remain calm and ensure your support person is with you. Time contractions. A bloody, mucous show may from your vagina may occur at this time. This can also occur a week before labour begins.

Note: contractions of the uterus that are not usually painful and often irregular are called Braxton Hicks and may occur throughout pregnancy.

Phone Birthing and Assessment Suites (or SMGP) before coming in to speak with a Midwife, who will support you in either staying at home longer or advise to come in.

Breaking of Membranes (waters)

If your membranes break (waters) put on a pad and check the colour of the amniotic fluid. Call Birthing and Assessment Suite and inform them.

The Midwife will ask questions about your condition:

- What time the membranes have broken and what colour the fluid is
- If the baby is moving and has there been change in movements
- What your contractions are like, how often and how long they last
- If there is any bright bleeding of the vagina

The Midwife will advise you when to come into Birthing and Assessment Suites.

Going to Hospital - Birthing and Assessment Suites

The midwife will advise to come in to Birthing and Assessment Suites if membranes have broken, if the discomfort/pain is too much to manage at home or if the midwife feels that assessment is required.

If you are with Southern Midwifery Group Practice, call your Midwife and she will instruct you further.

Birthing and assessment Suite

Birthing and assessment suite is located on the 3rd floor. The Midwife caring for you will answer any questions you may have and discuss any birthing preferences.

The midwife is there to support you throughout labour and to manage and monitor the progress.

Support during labour

The midwife will give support throughout labour but is advisable to have your partner and/or support person with you also.

Covid-19 has caused restrictions on how many people are allowed in rooms at any one time. Please phone Flinders Medical Centre for the current restrictions before you go into hospital to avoid disappointment (occasionally, if there are two support persons, one may be asked to leave due to room capacity constraints)

Care during labour

Throughout labour, the midwife will monitor temperature, blood pressure, pulse, contractions and the baby's heart rate.

If continuous monitoring is required, the midwife will discuss with you at the time.

Pain relief

During labour, a range of pain relief methods and options are available.

Common methods of relieving pain are:

- Being active
- Position changes
- Hot packs (please do not bring your own, these are provided)
- Water – shower or bath (a consent for the bath must be signed for in antenatal clinics prior)
- Nitrous oxide (gas)
- Tens machines
- Opiate injection (fentanyl)
- Subcutaneous water for injection in sacral pressure points
- Epidural anaesthesia

Induction of Labour

An Induction of Labour is when it is started artificially. This may be required if labour has not started naturally on its own or for other reasons explained by the doctor. The procedure will be fully explained to you prior to your admission to hospital.

Assisted delivery

Occasionally women require assistance at the end of labour to deliver their baby. This can be for a variety of reasons such as exhaustion or the baby heart rate suggests some distress. Types of assisted delivery that may occur are:

Episiotomy

It may be necessary to deliver the baby quickly or the opening may require to be enlarged in which an episiotomy is performed. This is a small incision in the perineum, between the vagina and the anus. This is usually a painless procedure, as local anesthetic is often used, and will be repaired post third stage with dissolvable stitches.

Forceps and Ventouse delivery

Forceps and ventouse may be required to assist the birth of the baby. If the delivery of baby requires either of these, the doctor will explain this with you at the time and a paediatrician will be present for the birth.

Caesarean section

A caesarean section is an operation that involves the delivery of the baby via a surgical cut through the abdomen and the uterus, usually near the top of the pubic area, however if baby requires delivery prematurely an abdominal incision higher may be required. Caesarean sections are performed with anaesthesia; either spinal, epidural top-up or general anaesthesia.

Approximately 30% of babies are born this way.

A caesarean can either be planned or an emergency. Reasons for a planned caesarean section will be discussed with you during antenatal appointments. An emergency caesarean section is performed if labour does not progress well.

After the birth of the baby

As soon as the baby is born, providing mother and baby are well, skin to skin contact is encouraged between mother and baby. If baby has been born by caesarean section, all attempts will be made to facilitate keeping the baby close to the mother as possible. If choosing to breast feed, this is encouraged straight after birth.

During admission or soon after birth, it will be asked if you consent to baby receiving Vitamin K. This is to prevent blood clotting disorders due to an underdeveloped clotting system. An information sheet regarding Vitamin K is located in the back of the orange South Australian Pregnancy Record.

Baby will be weighed and measured at this time.

Postnatal Ward and Care

After delivery of baby and when both mother and baby are stable, you will be transferred to the postnatal ward 4C or 4SMP. Depending on availability and care requirements the rooms are single or share rooms. Whilst it is not always possible, partners may be able to stay with you in the postnatal ward if you have a single room on the fold-out chair/bed. A donation for linen is appreciated.

The baby will “room-in” with the mother, so the mother can gain confidence in caring for and feeding her baby. Settling techniques and parenting skills will be demonstrated by the midwives. The baby will be referred to by the Mother’s surname to avoid confusion. Two name bands will be placed on the baby’s ankles for identification. Should a name band fall off, please alert the midwife to replace the identification band.

Once settled in the postnatal area, the midwife will spend time with the mother over the next few days (depending on how long your expected stay is) to educate on breast feeding and/or bottle feeding, bathing the baby, vaccinations the baby is recommended to have and prepare you for leaving the hospital.

Mother and baby will be assessed regularly each day to ensure wellbeing post birth.

Occasionally babies require extra care post birth and may be transferred to the Neonatal Unit. If baby requires care in the neonatal unit, every effort is made for the mother and/or father to be involved in the baby’s continuing care.

Visiting hours

Visiting hours are recommended between 3pm-8pm, however partners may be present outside of those times.

Covid-19 has brought in restrictions with visiting. Please contact the postnatal ward on 8204 4216 to receive the current visiting protocols.

If anyone is sick or unwell they will not be permitted in the hospital in any area.

Neonatal Unit

Sometimes babies require more care than the postnatal wards can give and are transferred to the Neonatal Unit, located on Level 3. This can be for a variety of reasons, which your doctor will explain and discuss with you. Parents are welcome to visit their baby; however strict visiting protocols guidelines are present.

Covid-19 has brought in strict restrictions with visiting in the neonatal area. Please contact the neonatal unit on 8204 5041 to receive the current visiting protocols.

Discharge from Hospital

The best place for mother and baby to be is in their own home environment. It is usual to stay in hospital one or two nights after baby is born, however women must stay in hospital for at least 4 hours post birth, and only if mother and baby are well will the hospital discharge them. Women who have a caesarean birth are expected to stay longer three or four nights.

Postnatal midwives will be preparing women for discharge and life at home. Discharge time is usually 10am. It is important to organise transport for this time.

Physiotherapy

Physiotherapists will see women who require extra care. Please discuss with your midwife.

Infant feeding

Breast Feeding

Flinders Medical Centre is an accredited Baby Friendly Health Initiative hospital. Breast feeding is recommended as the most beneficial way to feed your baby. Breast milk contains everything that your baby requires to grow and develop. Please refer to the "Advantages of Breast Feeding" education leaflet in the back of your orange South Australian Pregnancy Record.

It is important in your pregnancy that you obtain supportive maternity bras without wires, preventing blockages of the milk ducts. Try to avoid washing nipples with soap products when showering, as soap products will remove the protective natural oils and secretions.

Breast feeding is a learned technique which takes practice for both mother and baby. The midwives in the postnatal ward will educate and facilitate women to develop techniques for successful breastfeeding including correct positioning and attachment of the baby at the breast.

Artificial feeding

Women choosing to bottle feed will be supported by the midwives. Please bring a tin of your chosen formula for demonstration purposes. Formula and bottles will be provided whilst in hospital.

Registration Forms

All baby registration forms are online with Births, Deaths and Marriages. Parent packs are given out in Birthing and Assessment Suites.

Neonatal (Baby) examinations and tests

All babies will have a routine physical examination by a paediatrician prior to discharge. A routine Neonatal Screen Test is performed on all babies at 48 hours of life to test for rare metabolic diseases, that can be treated if detected early. This blood test requires a heel prick to gain blood. A hearing test on baby will be performed prior to discharged.

For further information, "Now that you have had your baby" book can be found on-line on the Flinders Medical Centre website under Women's and Children's Division.

Community Support

Maternity Outreach

Maternity Outreach Service will be offered to women and their baby within the defined Flinders Medical Centre catchment. A Midwife from Maternity Outreach will phone within 1-2 days after discharge. One or two visits will be offered as needed by a midwife. It is encouraged to make contact to organise follow-up with Child and Youth Health services and/ or your GP as well.

Post-natal day service

This is a free service for Medicare eligible women who might be finding difficulties in feeding or settling their baby. For more information or an appointment please phone the postnatal ward on 8204 4216.

Child and Youth Health Services (CYH)

Child and Youth Health Services aim to make contact with new mothers approximately 2-4 weeks after the birth of the baby, if consent given.

CYH can provide parenting support from birth to five years of age. Supports offered are breast feeding, developmental checks and support with any parenting concerns.

For earlier support or information please call CYH on 1300 733 606.

Australian Breast Feeding association (ABA)

This is a voluntary organisation that encourages and supports mothers wishing to breast feeding their babies. ANA has trained breast feeding counsellors who are available to assist mothers. For more information, please call the ABA on 8333 3276 or the ABA – breast feeding helpline on 1800 686 268.

Multiple Birth Association

The Multiple Birth association provide support and encouragement to parents of multiples of twins, triplets or more. For more information, please call the Multiple Birth association on 8342 2330. Flinders Medical Centre has a Multiple Birth Coordinator that can be contacted by calling Flinders Medical Centre on 8204 5511 and ask for pager 48033.

Postnatal Depression

Some women may experience “baby blues” shortly after the birth of their baby. Experiences may include feeling anxious and fearful. One in five mothers who feel these feelings do not improve over time and then can start feeling depression within the first 12 months. Depression is a treatable condition and often requires assistance to manage. Counselling and supportive therapy may all that is required, however some women may require medication.

Symptoms to be aware of are:

- Inability to sleep
- Decrease in interest, pleasure or motivation
- Irritability
- Constant fatigue
- Loss of appetite, weight changes
- Anxiety and panic feelings/thoughts
- Bad memories of childhood
- Disinterest in new baby

Please contact your GP should any of the feelings be experienced.

Alternatively phone one of these services for assistance:

- **Helen Mayo House**
(Family Unit Glenside)
8303 1183
(24-hour counselling line)
- **Domestic Violence Help Line**
1800 800 098
(24-hour counselling service)

- **Women’s Health State-wide**
1300 882 880 (metro callers)
1800 182 098 (country callers)
- **Emergency Department**
Flinders Medical Centre
8204 5511 and ask for the Emergency Department

Resources

Websites

Families – Services Australia

<http://www.servicesaustralia.gov.au/individuals/families>

(Centrelink Medicare Information)

Flinders Medical Centre

<http://www.sahealth.sa.gov.au/.../flinders+medical+centre>

Child and Youth Health Services

<https://www.cyh.com>

Australian Government Department of Health

<http://www.health.gov.au>

SA Health

<https://www.sahealth.sa.gov.au>

Parenting SA

<https://parenting.sa.gov.au>

Postnatal Depression

<https://www.beyondblue.org.au>

Perinatal Anxiety and Depression Australia

<https://www.panda.org.au>

Quitline

<https://www.health.gov.au/contacts/quitline>

Australian Institute of Social Relations

<https://www.socialrelations.edu.au>

(advice on relationship issues)

SA Multiple Birth Association

<https://www.multiplebirthsa.org.au>

Shine SA

<http://www.shinesa.org.au>

(sexual health/contraception)

Sids and Kids

<https://www.sidssa.org.au>

Australian Breast Feeding association (ABA)

<https://breastfeeding.asn.au>

Baby/Child restraint information

<https://www.mylicence.sa.gov.au/road-rules/seatbelts-and-child-restraints>

FLINDERS MEDICAL CENTRE IS A NON- SMOKING INSITUION AND PROMOTES HAND HYGIENE

Notes:

A series of horizontal dotted lines for writing notes.

For more information

**Flinders Women and Children
Flinders Medical Centre
Bedford Park
South Australia 5042
Telephone: 08 8204 5511
www.sahealth.sa.gov.au/fmc**



This document has been reviewed
and endorsed by consumers.



Interpreter



<https://creativecommons.org>

© Department for Health and Wellbeing,
Government of South Australia.
All rights reserved.
Printed September 2020



Government
of South Australia

Health
Southern Adelaide
Local Health Network