

# Abdominal Pain

## Patient discharge information sheet

This information sheet is designed to answer any questions you may have about your child having abdominal pain.

### What is Abdominal Pain?

Abdominal pain is pain or cramping anywhere in the abdomen (tummy, belly or stomach). Children often complain of abdominal pain. It is one of the most common reasons children see a doctor. Most cases of abdominal pain are not serious, and children often get better by themselves.

Pain can come from any structure in the abdomen including the stomach, bowel, kidneys, bladder and other organs. Even severe abdominal pain does not always mean your child is suffering from a serious illness. Severe pain can be associated with mild illnesses such as wind pain from drinking or eating too much or cramping from constipation or a viral infection. Some children get abdominal pain as a result of stress.

The exact cause of pain is often not found and will often get better by itself with simple pain relief and time.

There are, however, some serious illnesses that can start with mild abdominal pain.

### Signs and symptoms of abdominal pain

Abdominal pain can happen suddenly or develop slowly. Children often have other symptoms that are associated with the cause of the abdominal pain, such as:

- Vomiting
- Diarrhoea
- Fever

### What causes abdominal pain?

There are many things that can cause abdominal pain such as:

- Bowel (gut) problems – such as constipation or irritable bowel.
- Infections – such as gastroenteritis (which causes vomiting and diarrhoea/runny poo) or urine infections.
- Mesenteric adenitis – the lymph nodes in the abdomen commonly enlarge due to viral infections.
- Problems that may require surgery – such as appendicitis or a bowel obstruction.
- Period pain – monthly pain can occur before or during a menstrual period.

- Food related – too much food, food poisoning or food allergies and intolerances.
- Some children get abdominal pain as a result of stress or anxiety.
- Sometimes there is no identifiable cause for the abdominal pain.
- The causes of abdominal pain can be hard to determine. Sometimes the cause becomes more obvious with time, and then doctors can work out the best treatment.

### When to see a doctor

Many children with abdominal pain get better quickly without any treatment and there is no need to see a doctor. If your child's pain or problems persist for more than 24 hours, or you are worried about your child, take them to your GP.

Take your child to the GP or hospital as soon as possible if your child:

- has severe pain (despite pain medication) or the pain has moved
- has pain that returns frequently and regularly
- does not want to move
- has a fever (temperature over 38 degrees)
- is pale, sweaty, lethargic (hard to wake) and unwell
- is refusing to drink fluids
- is vomiting for more than 24 hours and not keeping fluids down, or their vomit is green in colour
- has blood in their vomit or faeces (poo)
- has problems passing urine (doing a wee)
- is a baby, and has less than four wet nappies a day
- has pain or lumps in their groin or testicles in a boy
- has a skin rash which is sore or painful
- has had a recent injury (for example, falling onto bike handlebars).

### Treatment for abdominal pain

Treatment may be as simple as going home to rest, drink fluids and eat a bland diet. At other times, your child may be admitted to hospital or may need an operation (surgery). Sometimes tests are needed to help work out the cause of the pain. These may include:

- blood tests
- a urine test
- a stool (poo) sample
- X-rays of the abdomen
- ultrasound.

## Repeated attacks of abdominal pain

Some children get repeated attacks of abdominal pain, which can be very worrying for parents. Often no health problem can be found. Children may have abdominal pain when they are worried about themselves or people around them.

Think about whether there is anything that is upsetting your child at home, at school, kindergarten or with friends. See your GP for advice. Your child may need a referral to a paediatrician, gastroenterologist (a doctor who specialises in stomach problems) or psychologist.

## Care at home

Here are some general ways to ease your child's pain:

- Help your child drink their usual amount of fluids. Getting your child to drink is important as it prevents dehydration (loss of water).
- If your child is hungry, let them eat what they want or offer bland foods such as crackers, rice, bananas or toast. Do not force your child to eat if they feel unwell. They will start eating again when they feel better.
- Encourage sitting on the toilet. Sometimes doing a poo helps to ease the pain.
- Rubbing a child's tummy or having a distraction, such as reading a book, can sometimes ease the pain.
- Give paracetamol or ibuprofen if your child is in pain or is miserable. If ibuprofen causes a stomach upset, offer your child some food or milk.

## Key points to remember

- Many children get abdominal pain and most get better by themselves.
- Make sure your child drinks plenty of fluids to prevent dehydration.
- Often no cause can be found, and sometimes a cause becomes more obvious with time.
- If your child has abdominal pain and looks unwell, take your child to your GP or local hospital as soon as possible

## Common questions our doctors are asked

### How can I tell if my child has appendicitis?

Appendicitis can be difficult for doctors to diagnose, but a sign that your child may have appendicitis is that they have severe pain starting around their belly button and moving to the right side of their abdomen. Most children with appendicitis will be very reluctant to move. See your GP if you are worried.

### My child has been diagnosed with mesenteric adenitis. What does this mean?

Mesenteric adenitis occurs when the lymph nodes in the abdomen enlarge in response to an infection – most commonly a viral infection. This results in stomach pain. For further information see Mesenteric Adenitis Fact Sheet.

### Why is it so difficult to work out the cause of my child's ongoing stomach aches?

Stomach aches are difficult to diagnose in all ages. Children differ in their ability to describe the type, severity and location of their pain, which can make this process even harder. Many problems from the chest down to the groin may be interpreted by children as stomach aches, making it very difficult to find out the true cause. Your child's doctor will examine and investigate your child in order to rule out anything serious, while trying to find the underlying cause.

### The doctor says my child has abdominal migraine. What is this?

As the name suggests, abdominal migraine is a migraine experienced in the abdomen instead of the head. A child with abdominal migraine will often have tummy pain along with nausea/vomiting, loss of appetite and pale skin. There is no headache involved and the child is well between episodes. There is still quite a lot that is unknown about abdominal migraine, but the risk factors and triggers are thought to be similar to traditional migraines (e.g. having a family member with migraines, being stressed or overtired, chemicals in food).

## For more information

See your family doctor  
 For 24 hour health advice call - Health Direct Australia 1800 022 222  
 Parent Helpline 1300 364 100  
 Local emergency department

Adapted from Royal Childrens Melbourne (Abdominal Pain – acute) UpToDate: abdominal Pain (the basics) Health direct (abdominal pain in children) Childrens hospital westmead (abdominal Pain)

## For more information

Womens and Childrens Division

Flinders Medical Centre

Flinders Drive. Bedford Park, South Australia 5042

Telephone: 08 8204 5511

[www.sahealth.sa.gov.au/fmc](http://www.sahealth.sa.gov.au/fmc)

© Department for Health and Wellbeing, Government of South Australia.

Reviewed August 2023. Next review August 2026.



This document has been reviewed and endorsed by consumers.



Government of South Australia

Health

Southern Adelaide  
Local Health Network