

Clozapine Temperature Monitoring

Important information on how to manage a high temperature or fever during the first 28 days of treatment with clozapine.

Record extra temperature or your notes here:

| Start date: | | | | | | |
|-------------|----|----|----|----|----|----|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| 8 | 9 | 10 | 11 | 12 | 13 | 14 |
| 15 | 16 | 17 | 18 | 19 | 20 | 21 |
| 22 | 23 | 24 | 25 | 26 | 27 | 28 |
| | | | | | | |





Clozapine Temperature Monitoring

Important information on how to manage a high temperature or fever during the first 28 days of treatment with clozapine.

Record extra temperature or your notes here:

| Start date: | | | | | | |
|-------------|----|----|----|----|----|----|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| 8 | 9 | 10 | 11 | 12 | 13 | 14 |
| 15 | 16 | 17 | 18 | 19 | 20 | 21 |
| 22 | 23 | 24 | 25 | 26 | 27 | 28 |
| End date: | | | | | | |

of South Australia

Clozapine Temperature Monitoring

Important information on how to manage a high temperature or fever during the first 28 days of treatment with clozapine.

Record extra temperature or your notes here:

| Start date: | | | | | | |
|-------------|----|----|----|----|----|----|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| 8 | 9 | 10 | 11 | 12 | 13 | 14 |
| 15 | 16 | 17 | 18 | 19 | 20 | 21 |
| 22 | 23 | 24 | 25 | 26 | 27 | 28 |
| End date: | | | | | | |



Why should I check my temperature?

- A raised temperature is common in the first 28 days of clozapine treatment, and usually means that it is starting to work.
- In rare cases a high temperature or feeling feverish can be signs of a severe blood infection in the body or a heart condition.
- It is important to check your temperature every day to safeguard your wellbeing and help detect problems early on.
- If you don't have your own thermometer, you will be given one like this to keep:



What do I need to do?

- Check your temperature under your tongue or your armpit at approximately the same time every day, for the first 28 days
- If placing under your tounge make sure you have not had a hot or cold drink for at least 20 minutes beforehand
- Record your temperature in the table overleaf
- Clean thermometer after each use and store safely
- Note: You can also check your temperature any other time you feel unwell

High temperature instructions

- If your temperature is 38 degrees or more
 - Check your temperature again after 15 minutes
 - Telephone your GP or Clozapine Coordinator who will advise you what to do
 - If it is a weekend or public holiday go to a hospital Emergency Department.
 Tell the staff that you are on clozapine, and that your temperature is raised.
 Some extra blood tests and heart checks may be needed and/or changes in treatment.

Why should I check my temperature?

- A raised temperature is common in the first 28 days of clozapine treatment, and usually means that it is starting to work.
- In rare cases a high temperature or feeling feverish can be signs of a severe blood infection in the body or a heart condition.
- It is important to check your temperature every day to safeguard your wellbeing and help detect problems early on.
- If you don't have your own thermometer, you will be given one like this to keep:



What do I need to do?

- Check your temperature under your tongue or your armpit at approximately the same time every day, for the first 28 days
- If placing under your tounge make sure you have not had a hot or cold drink for at least 20 minutes beforehand
- Record your temperature in the table overleaf
- Clean thermometer after each use and store safely
- Note: You can also check your temperature any other time you feel unwell

High temperature instructions

- If your temperature is 38 degrees or more
 - Check your temperature again after 15 minutes
 - Telephone your GP or Clozapine Coordinator who will advise you what to do
 - If it is a weekend or public holiday go to a hospital Emergency Department.
 Tell the staff that you are on clozapine, and that your temperature is raised.
 Some extra blood tests and heart checks may be needed and/or changes in treatment.

Why should I check my temperature?

- A raised temperature is common in the first 28 days of clozapine treatment, and usually means that it is starting to work.
- In rare cases a high temperature or feeling feverish can be signs of a severe blood infection in the body or a heart condition.
- It is important to check your temperature every day to safeguard your wellbeing and help detect problems early on.
- If you don't have your own thermometer, you will be given one like this to keep:



What do I need to do?

- Check your temperature under your tongue or your armpit at approximately the same time every day, for the first 28 days
- If placing under your tounge make sure you have not had a hot or cold drink for at least 20 minutes beforehand
- Record your temperature in the table overleaf
- Clean thermometer after each use and store safely
- Note: You can also check your temperature any other time you feel unwell

High temperature instructions

- If your temperature is 38 degrees or more
 - Check your temperature again after 15 minutes
 - Telephone your GP or Clozapine Coordinator who will advise you what to do
 - If it is a weekend or public holiday go to a hospital Emergency Department.
 Tell the staff that you are on clozapine, and that your temperature is raised.
 Some extra blood tests and heart checks may be needed and/or changes in treatment.