South Australian Neonatal Medication Guidelines

Pancuronium

(SAS) 4mg/2mL injection © Department for Health and Wellbeing, Government of South Australia. All rights reserved.

This guideline provides advice of a general nature. This statewide guideline has been prepared to promote and facilitate standardisation and consistency of practice, using a multidisciplinary approach. The guideline is based on a review of published evidence and expert opinion.

Information in this statewide guideline is current at the time of publication.

SA Health does not accept responsibility for the quality or accuracy of material on websites linked from this site and does not sponsor, approve or endorse materials on such links.

Health practitioners in the South Australian public health sector are expected to review specific details of each patient and professionally assess the applicability of the relevant guideline to that clinical situation.

If for good clinical reasons, a decision is made to depart from the guideline, the responsible clinician must document in the patient's medical record, the decision made, by whom, and detailed reasons for the departure from the guideline.

This statewide guideline does not address all the elements of clinical practice and assumes that the individual clinicians are responsible for discussing care with consumers in an environment that is culturally appropriate and which enables respectful confidential discussion. This includes:

- The use of interpreter services where necessary,
- Advising consumers of their choice and ensuring informed consent is obtained,
- Providing care within scope of practice, meeting all legislative requirements and maintaining standards of professional conduct, and
- Documenting all care in accordance with mandatory and local requirements

This is a High Risk Medication 🛝



Only muscle-relax a neonate if confident that the airway can be maintained and that hand ventilation can be provided

Dose and Indications

Skeletal muscle paralysis in patients with assisted ventilation

Intravenous

Weight	Dose*	
<1kg	0.1mg flat dose	
>1kg	0.1mg/kg/dose	

Repeat one to two hourly if needed. Adjust dose as needed based on duration of paralysis

Preparation and Administration

Intravenous

Dilute 1mL of the 2mg/mL pancuronium solution with 1mL of compatible fluid (to a total volume of 2mL). The resulting solution contains 1mg/mL pancuronium.

	0.1mg				
Volume	0.1mL	0.2mL	0.3mL	0.4mL	0.5mL

Administer as a rapid intravenous push over a few seconds

Discard remaining solution.

This product is not registered in Australia and is accessed under the Special Access Scheme (SAS). Please complete relevant TGA paperwork.

SA Health

INFORMAL COPY WHEN PRINTED Page 1 of 2

^{*}Dose has been rounded to prevent tenfold errors in babies less than 1kg

Pancuronium (SAS) 4mg/2mL injection

Compatible Fluids

Glucose 5%, glucose / sodium chloride solutions, sodium chloride 0.9%

Adverse Effects

Common

Hypertension, tachycardia, prolonged paralysis

Rare

Anaphylactic reactions

Note: Hypoxaemia may occur because of inadequate mechanical ventilation and deterioration in pulmonary mechanics

Monitoring

- > Continuous cardio-respiratory and pulse oximetry
- > Blood pressure

Practice Points

- > Use only if patient is on assisted ventilation.
- > Provide eye protection as needed and instil lubricating eye drops every 2 hours
- > To reverse the effects of pancuronium; use neostigmine with atropine
- > The manufacturer recommends that pancuronium bromide not be mixed with other drugs in a syringe as possible changes in pH may result in precipitation.

Document Ownership & History

Developed by: SA Maternal, Neonatal & Gynaecology Community of Practice

Contact: Health.NeoMed@sa.gov.au

Endorsed by: SA Health Safety and Quality Strategic Governance Committee

Next review due: 24/04/2023

ISBN number: 978-1-74243-966-2

PDS reference: CG048

Policy history: Is this a new policy (V1)? **N**

Does this policy amend or update and existing policy? Y

If so, which version? V1

Does this policy replace another policy with a different title?

If so, which policy (title)?

Approval Date	Version	Who approved New/Revised Version	Reason for Change
9/04/21	V2.1	Deputy CE, Commissioning and Performance, SA Department for Health and Wellbeing	Product change
24/04/18	V2	SA Health Safety and Quality Strategic Governance Committee	Formally reviewed in line with 5 year scheduled timeline for review.
1/12/12	V1	SA Health Safety and Quality Strategic Governance Committee	Original SA Health Safety and Quality Strategic Governance Committee approved version.