Port Pirie Blood Lead Levels

Analysis of blood lead levels for 2021 (1 January – 31 December 2021)



Background

Situated 230 km north of Adelaide in South Australia, Port Pirie is the location of one of the world's largest lead and multi metal smelters.

The smelter has been in continuous operation since 1889. Over time, airborne lead-contaminated dust produced during smelter operations has contaminated the local environment. Even small amounts of lead can be toxic when ingested or inhaled. Notwithstanding ongoing improvements to practices that reduce dust, lead dust emissions from the smelter, together with legacy lead from past operations, are a persistent source of exposure for the Port Pirie community.

SA Health has provided voluntary blood lead screening through the Port Pirie Lead Implementation Program that has been delivered by the Port Pirie Environmental Health Centre since 1984. This Program monitors and helps the local community reduce the amount of lead that children absorb by providing lead exposure-reduction advice, education and interventions for families. This paper reports the analysis of blood lead levels of Port Pirie children under 5 years of age (0-1824 days of age i.e. up to the date of a child's fifth birthday) and pregnant women that have been tested each year since 2012. Reports summarising data from the blood testing program for years before 2012 are available on request.

Australian lead guidelines

The National Health and Medical Research Council (NHMRC) provides health advice and guidelines for the Australian community, governments and health professionals. One of the NHMRC's tasks is to advise the Australian community about lead exposure and the health effects of lead and how they can be managed.

Guideline for investigating lead exposure

The NHMRC recommends that if a person has a blood lead level greater than 5 micrograms per decilitre (μ g/dL), the source of lead exposure should be investigated and reduced, particularly if the person is a child or pregnant woman. See the <u>Frequently Asked Questions: NHMRC Review of blood</u> lead level guidelines.

The NHMRC advises it is well established that exposure to lead at blood lead levels above 10 μ g/dL can have harmful effects on body functions and organs in both adults and children. Research now shows that blood lead levels below 10 μ g/dL may also be associated with some health effects (<u>NHMRC 2015</u>, Evidence on the effects of lead on human health). At this stage, the NHMRC has concluded that the evidence is not strong enough to show that lead is the cause of these effects.

In line with NHMRC's guidance, the blood lead levels of children living in Port Pirie are reported against the exposure investigation level of 5 μ g/dL. This is in addition to the target level of 10 μ g/dL that has been reported each year since 2000 and was the target level of the Targeted Lead Abatement Program when it commenced in 2014.

Blood lead levels of Port Pirie children under 5 years of age

Blood lead levels above 5 micrograms per decilitre

In 2021, 385 children had a reported result above the NHMRC blood lead exposure investigation level of 5 μ g/dL (Table 1 and Figure 1).

Blood test results indicate that 64.5% of the children tested in 2021 had blood lead levels above 5 μ g/dL. This is a 3.7% decrease compared to 2020.



These percentages cannot be reported or interpreted to represent the proportion of the Port Pirie population above 5 μ g/dL because not every child living in Port Pirie was tested in 2021, therefore their blood results do not appear in this analysis. It can be said with a high degree of confidence (95%) that the proportion of the entire population with blood lead levels above 5 μ g/dL would fall between the lower limit of 62.9% and the upper limit of 66.1% (the 95% confidence interval) if every child under 5 years of age living in Port Pirie was tested (Table 1 and Figure 1). Population estimates used to calculate these limits are based on Estimated Resident Population (ERP) data for Port Pirie provided by the Australian Bureau of Statistics (ABS).

The 95% confidence intervals that contain the population proportions may vary between reports because they are adjusted from year to year for the estimated population size.

Table 1: Percentage of children under 5 years of age (0-4yrs) tested with blood lead levels above the Lead Exposure Investigation Level >5 μ g/dL and the lower limit and upper limit (95% confidence interval) for each calendar year 2012-2021.

	ERP	Total		>5 µ <u>(</u>	g/dL	
Year	estimated number of children under 5 years of age	number of children tested	n	Lower limit	%	Upper limit
2012	858	555	276	47.3	49.7	52.2
2013	835	569	303	50.9	53.3	55.6
2014	809	560	284	48.4	50.7	53.0
2015	773	556	264	45.3	47.5	49.7
2016	786	571	279	46.7	48.9	51.0
2017	747	550	288	50.2	52.4	54.5
2018	755	515	296	55.0	57.5	59.9
2019	769	564	359	61.6	63.7	65.7
2020	757	553	377	66.1	68.2	70.2
2021	720	597	385	62.9	64.5	66.1

Blood lead levels measured at the time of children's second birthdays (2-year-old children) are considered to be robust indicators of lead exposure for the whole population. This is also the age of children who have the highest level of participation in the blood testing program so results for this age group give the best representation of trend.

In 2021, 63 children tested at their second birthday had a reported result above the NHMRC blood lead investigation level of 5 μ g/dL (Table 2 and Figure 1).

Blood test results indicate that 75.9% of children tested at their second birthday in 2021 had blood lead levels above 5 μ g/dL. This is a 5.8% increase compared to 2020.




Table 2: Percentage of children tested at their second birthday (2-year-old children) with blood lead levels above the Lead Exposure Investigation Level >5 μ g/dL and the lower limit and upper limit (95% confidence interval) for each calendar year 2012-2021.

	ERP	Total		>5 µg	g/dL	
Year	estimated number of children at their second birthday	number of 2-year old children tested	n	Lower limit	%	Upper limit
2012	155	107	64	54.5	59.8	64.9
2013	166	118	71	55.3	60.2	64.9
2014	186	135	67	45.2	49.6	54.1
2015	172	119	66	50.4	55.5	60.4
2016	171	113	61	48.5	54.0	59.3
2017	148	94	52	49.1	55.3	61.4
2018	165	93	58	55.6	62.4	68.7
2019	133	90	64	65.4	71.1	76.2
2020	171	97	68	63.7	70.1	75.8
2021	147	83	63	69.2	75.9	81.5

CAVEAT: there is no accurate ABS estimate for the number of children who turned 2 years of age in each calendar year, therefore an estimate has been extrapolated from ABS birth data for Port Pirie in 2 years prior to the reporting year to derive the lower and upper limits for this age group.

Figure 1: Percentage of tested children with blood lead levels >5 μ g/dL for children under 5 years of age (0-4yrs) and children at their second birthday (2-year-old children) for each calendar year 2012-2021. Bars indicate the lower limit and upper limit of the 95% confidence interval.





Results in each category of blood lead level

Blood test results indicate that 41% of children tested in 2021 had blood lead levels between 5 μ g/dL and 10 μ g/dL, 16.1% between 10 μ g/dL and 15 μ g/dL and 6.4% between 15 μ g/dL and 20 μ g/dL (Table 3 and Figure 2).

The number of children with blood lead levels equal to or exceeding 20 μ g/dL (6 children) has decreased compared to the number reported for 2020 (16 children were reported in the 2020 annual report. However, a routine audit identified a subsequent confirmational test of one of these children returned after finalising the report with a level below 20 μ g/dL which means that overall, 15 children had blood lead levels equal to or exceeding 20 μ g/dL measured in 2020).

Table 3: Percentage of children under 5 years of age (0-4yrs) tested in each blood lead level category
for each calendar year 2012-2021.

Year	Number and percentage	Children ≤5 μg/dL	Children >5 and <10 μg/dL	Children ≥10 and <15 µg/dL	Children ≥15 and <20 μg/dL	Children ≥20 µg/dL	Total number of children tested
2012	n	279	137	87	38	14	555
2012	%	50.3%	24.7%	15.7%	6.8%	2.5%	000
2013	n	266	174	88	26	15	569
2010	%	46.7%	30.6%	15.5%	4.6%	2.6%	303
2014	n	276	168	80	29	7	560
2014	%	9.3%	30%	14.3%	5.2%	1.3%	500
2015	n	292	161	77	18	8	556
2015	%	52.5%	29%	13.8%	3.2%	1.4%	330
2016	n	292	155	91	23	10	571
2010	%	51.1%	27.1%	15.9%	4%	1.8%	571
2017	n	262	165	74	43	6	550
2017	%	47.6%	30%	13.5%	7.8%	1.1%	330
2018	n	219	152	91	37	16	515
2010	%	42.5%	29.5%	17.7%	7.2%	3.1%	515
2019	n	205	211	99	36	13	564
2019	%	36.3%	37.4%	17.6%	6.4%	2.3%	004
2020	n	176	221	99	42	15 (16)*	553
2020	%	31.8%	40%	17.9%	7.6%	2.7%	222
2024	n	212	245	96	38	6	507
2021	%	35.5%	41%	16.1%	6.4%	1%	597

* 16 children were reported in the 2020 annual report to have blood lead levels equal to or exceeding 20 μg/dL. However, a routine audit identified a subsequent confirmational test of one of these childrem

returned after finalising the report with a level below 20 μ g/dL which means that overall, 15 children had blood lead levels equal to or exceeding 20 μ g/dL measured in 2020.




Figure 2: Percentage of children under 5 years of age (0-4yrs) tested in each blood lead level category for each calendar year 2012-2021

Blood lead levels equal to or above 10 micrograms per decilitre

Blood tests indicate that 23.5% of children tested had blood lead levels equal to or above 10 μ g/dL. This is a 4.7% decrease compared to 2020 (Table 4).

Table 4: Percentage of children aged under 5 years of age (0-4 yrs) tested with blood lead levels \geq 10 µg/dL and the lower limit and upper limit (95% confidence interval) for each calendar year 2012-2021.

	ERP estimated		≥10 µg/dL				
Year	number of children under 5 years of age	Total number of children tested	n	Lower limit	%	Upper limit	
2012	858	555	139	23.0	25.0	27.3	
2013	835	569	129	20.8	22.7	24.7	
2014	809	560	116	18.9	20.7	22.6	
2015	773	556	103	16.9	18.5	20.3	
2016	786	571	124	20.0	21.7	23.5	
2017	747	550	123	20.6	22.4	24.2	
2018	755	515	144	25.8	28.0	30.2	
2019	769	564	148	24.4	26.2	28.2	
2020	757	553	156	26.3	28.2	30.2	
2021	720	597	140	22.1	23.5	24.9	



Blood tests indicate that 31.3% of children tested at their second birthdays had blood lead levels equal to or above 10 μ g/dL. This is a 1.7% decrease compared to 2020 (Table 5).


Table 5: Percentage of children tested at their second birthday (2-year-old children) with blood lead levels \geq 10 µg/dL and the lower limit and upper limit (95% confidence interval) for each calendar year 2012-2021.

	ERP	Total	≥10 µg/dL					
Year	estimated number of children at their second birthday	number of 2-year old children tested	n	Lower limit	%	Upper limit		
2012	155	107	32	25.3	29.9	35.0		
2013	166	118	27	19.0	22.9	27.3		
2014	186	135	25	15.3	18.5	22.2		
2015	172	119	30	21.1	25.2	29.8		
2016	171	113	22	15.5	19.5	24.1		
2017	148	94	18	14.7	19.1	24.5		
2018	165	93	27	23.3	29.0	35.6		
2019	133	90	30	28.0	33.3	39.2		
2020	171	97	32	27.1	33.0	39.5		
2021	147	83	26	25.1	31.3	38.4		

CAVEAT: there is no accurate ABS estimate for the number of children who turned 2 years of age in each calendar year, therefore an estimate has been extrapolated from ABS birth data for Port Pirie in 2 years prior to the reporting year to derive the lower and upper limits for this age group.



Figure 3: Percentage of tested children with blood lead levels $\geq 10 \mu g/dL$ for children under 5 years of age (0-4yrs) and children at their second birthday (2-year-old children) for each calendar year 2012-2021. Bars indicate the lower limit and upper limit of the 95% confidence interval.



Average blood lead level for children tested on their second birthday

Blood tests indicate that the average blood lead level (geometric mean) of children tested on their second birthday (2-year-old children) in 2021 was 7.3 μ g/dL (Table 6). This average has not changed compared to 2020.

Table 6: Geometric mean blood lead levels for children tested on their second birthday (2-year-old children) for each calendar year 2012-21.

Year	Number of 2- year-old children tested	Geometric mean of 2-year old children tested (µg/dL)
2012	107	6.3
2013	118	6.1
2014	135	5.5
2015	119	5.9
2016	113	5.5
2017	94	5.6
2018	93	5.8
2019	90	7.2
2020	97	7.3
2021	83	7.3



Blood lead levels of pregnant women living in Port Pirie

Blood tests indicate that the average blood lead level (geometric mean) of pregnant women living in Port Pirie who were tested in 2021 was 1.4 μ g/dL (Table 7). This average has decreased by 0.3 μ g/dL compared to 2020. This measure represents the blood lead level that a child has at birth.

Table 7: Geometric mean blood lead levels for pregnant women tested (during pregnancy or immediately postpartum) for each calendar year 2012-21.

Year	Number of pregnant women tested	Geometric mean of pregnant women tested (µg/dL)
2012	195	1.6
2013	175	1.5
2014	174	1.4
2015	196	1.5
2016	183	1.3
2017	143	1.4
2018	181	1.4
2019	158	1.8
2020	173	1.7
2021	170	1.4

Population profile

Population estimates indicate a decreasing number of children under 5 years of age living in Port Pirie (Table 8). Figure 4 represents this declining trend over the past 10 years from the ERP provided by the ABS that has been derived from census data.

This report uses a population estimate for 2021 that is a linear extrapolation from ERP of children under 5 years of age (0-4 years) living in Port Pirie (SA2 location 405031122) over the past 10 years. The ERP is updated by the ABS between censuses and completely revised each time a census is conducted. The ERPs used in this analysis are based on data for Port Pirie updated by the ABS on 28 August 2021 using migration and birth rates in addition to 2011 and 2016 census data.

The ERP is the most reliable estimate of the population between census years because it accounts for changes in population such as births, deaths, and the movement of residents in and out of the region in addition to the most recent census data. While the ABS states that the ERP is the most accurate estimate of the population, there is inherent inaccuracy involved in estimating population, with the largest margin of error recorded with small population sizes such as the population of children under 5 years of age living in Port Pirie. Any minor change in determinants of population size such as migration and birth rate can cause a substantial change in population estimates as a result of the small population size.

Although there is no accurate ABS estimate for the number of children who turned 2 years of age (have their second birthday) in each calendar year, an estimate has been extrapolated from ABS birth data for Port Pirie in 2 years prior to the reporting year to derive the lower and upper limits for this age group in this report.




Table 8: ERP data provided by ABS for children under 5 years of age living in Port Pirie Statistical Area Level 2 (SA2) for each calendar year 2012-2020 and extrapolated population estimate for 2021.

Age	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Estimated number of children under 5 years of age*	858	835	809	773	786	747	755	769	757	720
Estimated number of children at their second birthday [‡]	155	166	186	172	171	148	165	133	171	147

*Population estimates for children under 5 years of age living in Port Pirie SA2 (location code 405031122) each calendar year 2012-2021 are taken directly from ABS ERP data, except for 2021 which is a population estimate extrapolated from a linear model fitted to ERP data from the previous 10 years.

[‡]Population estimates for children at their second birthday are made directly from ABS births data for Port Pirie SA2 in the calendar year 2 years prior – e.g. number of children who turned 2 years of age in 2021 = number of births in 2019.

Figure 4: ERP data provided by ABS for children under 5 years of age living in Port Pirie Statistical Area Level 2 (SA2) for each calendar year 2012-2020 and extrapolated population estimate for 2021



Source: Linear extrapolation from the Australian Bureau of Statistics Estimated Resident Population 2012-2020 for Port Pirie (SA2 location code <u>405031122</u>).

Analysis methods

Considerations when comparing current results with historical data and reports

Multiple methods of analysis and reporting that have been undertaken since the Port Pirie lead program commenced in the 1980s that impact on comparison of blood lead trends over the 10-year period reported here, or comparison of these results to historical reports. Consideration should be given to the following:



- > Blood testing service providers for the Port Pirie community blood lead screening program changed on 1 January 2019, allowing for more advanced analytical methods to be utilised. This change should be considered when comparing data from 2019 onwards to previous years.
- > From, 2007 to June 2021 results were reported with surrogates where surrogate data (the mother's blood lead levels) was used to represent a child's blood lead level at birth and was included until either the child's first test at 6 months of age replaced it or it was removed if the child did not have any test by 9 months of age. From July 2021, results are reported separately either from pregnant women tested or from children under 5 years of age (0-4 years) with their first test usually at 6 months of age tested in the reporting period. Reports from July 2021 onwards should not be compared with reports from previous years containing surrogates
- > Routine data audit occurs when each analysis is undertaken and will continue to identify small numbers of data collection and entry anomalies (e.g. clients reporting incorrect family details). Correction of these anomalies can result in a minor variation in the number of test results included in the dataset compared to reports prepared in previous years. But this correction does not have significant impact on reported measures.

Blood lead data profile

Children included in this report were under 5 years of age (0-1824 days of age which is the time up to the date of a child's fifth birthday) and lived in Port Pirie at the time the blood test was taken. This is their age at the time the test was taken, not their current age. If more than one blood test was taken for a child in the reporting period, the child's most recent test result was used in this analysis. Thus, this analysis can include a child's test result that may have been collected any time up to 364 days prior to the preparation of this report. For this reason, a child's most recent test result may not represent the child's current blood lead level.

All blood samples included in this report are analysed by Australian laboratories accredited for blood lead analysis by the National Association of Testing Authorities (NATA Australia).

Sampling bias in blood lead data

These results cannot be interpreted to represent the entire population because test results for every child living in Port Pirie are not in every report because:

- > Although there is a high level of community participation in the blood lead screening program, the program is voluntary and therefore parents have the right to choose not to have their child tested. For this reason, there is no ethical way to mandate testing to test every child.
- Some children have a history of consistently low blood lead levels and have been assessed to remain at a low risk of lead exposure and developing lead-related health effects. These children will not benefit from ongoing testing beyond 3 years of age as part of their lead exposure management strategy. However, children at high risk of lead exposure are always encouraged to be regularly tested.
- Results pending re-test for confirmation are not included in this dataset but will appear in the next report.

These reports are considered very reliable for reporting the proportion of the population with blood lead levels above 10 micrograms per decilitre (the children at highest risk). However, children with low blood lead levels are under-represented i.e. there is inherent bias in this analysis (for the reasons given above). To help address some of this sampling bias, a statistical range is reported (the 95% confidence interval). It can be said with a high degree of confidence (95%), that the proportion of the entire population with blood lead levels above 5 µg/dL or equal to or above 10 µg/dL would fall



between the lower limit and the upper limit (the 95% confidence interval) of this range if every child under 5 years of age living in Port Pirie was tested.

Further information about the analysis methods used in this paper is available at <u>Frequently Asked</u> Questions Testing and Reporting Port Pirie Children's blood lead levels.

This report prepared by: Scientific Services Branch, Health Protection and Licensing Services

February 2022 Doc. no. A3322138 File no. 2021-05423

For more information

Scientific Services Branch Health Protection and Licensing Services, SA Health PO Box 6, Rundle Mall Adelaide SA 5000 Telephone: 08 8226 7100 www.sahealth.sa.gov.au

Public-I1-A1





