Fact sheet

Central Adelaide Ophthalmology Service

Clinical Information Sheet

Clinical Condition	DIABETIC EYE DISEASE
Eligibility	Retinal assessment including VA and refraction with an Optometrist or Ophthalmologist. Referral category dependent on clinical need. Clinical need is defined by NHMRC Guidelines: http://www.nhmrc.gov.au/guidelines/publications/di15
Priority and how to access services	All people with diabetes require regular retinopathy screening by an ophthalmologist or optometrist using ophthalmoscopy or photography. Type 1: Refer 5 years after diagnosis - non-urgent (cat 3) Type 2: Refer at diagnosis - non-urgent (cat 3) except if retinopathy present or patient pregnant - semi urgent (cat 2). Diabetes with sudden vision loss - Emergency
Differential Diagnoses	Retinal vein occlusion Hypertensive eye disease
Information required with referral	Optometrist/Ophthalmologist report including VA, refraction and impact of symptoms. Level of visual impairment (recreational, educational, occupational, driving)
Investigations required with referral	Blood sugar level, HbA1C, blood pressure, cholesterol (GP) OCT macula if available (optometry/ophthalmology)
Pre-Referral management strategies (information required with referral)	Ensure tight BSL, BP and cholesterol control
Discharge Criteria/information	Patients require regular follow up guided by the stage of their disease
Fact sheets	http://www.nhmrc.gov.au/guidelines/publications/di15 http://www.ranzco.edu/images/NEW_SITE/Eye_Health_Information /opa_ranzco_diabetic_retinopathy.pdf

For more information

Central Adelaide Ophthalmology Service Royal Adelaide Hospital, North Terrace ADELAIDE Telephone: 08 8222 4000 The Queen Elizabeth Hospital, 28 Woodville Road, WOODVILLE Telephone: 08 8222 6000

