Fact sheet

Central Adelaide Orthopaedics – Hip & Knee Service

Clinical Information Sheet

Clinical Condition	Baker's Cyst
Clinical Condition	Baker's Cyst
Eligibility	Swelling behind the knee
	Stiffness in the knee
	Knee pain
Priority	Not Eligible: patients with a diagnosis of bakers cyst will not be offered an
	appointment
	Assess patient for underlying condition, such as arthritis or a cartilage tear for example
Differential	Popliteal artery aneurysm
Diagnoses	Tumour
ha farma a filoro	Definite the flavorist flat and the flavorist transfer
Information required with	Patients with a diagnosis of bakers cyst will not be offered an appointment Assess patient for underlying condition, such as arthritis or soft tissue damage
referral	Assess patient for underlying condition, such as artifitis or soit tissue damage
Investigations	All patients should be assessed by plain x-ray as a basis to establish bony
required with	structure and eliminate osteoarthritis as an underlying cause
referral	X-ray: AP, Lateral knee & patella skyline view
	Consider Ultrasound or MRI to exclude other pathologies
	Upon attendance to appointment, patient will be required to bring plain x-rays
	(views mentioned above) from within the previous 6 months to establish
	current bony structure
Pre-Referral	Ice the area
management	Compression bandaging
strategies (include	Physiotherapy for knee strengthening exercises
with referral)	Use of simple analgesia as tolerated including a regular paracetamol product (e.g. Panadol® Osteo) and oral NSAIDs if tolerated
	Consider hydrotherapy, swimming or cycling for a low-impact exercise
	alternative
	Use of viscosupplementation (e.g. high dose fish oil)
	Use of mobility aids (e.g. walking stick or crutches)
	Weight loss measures
Discharge	Once underlying cause established, refer patient for underlying cause
Criteria/information	, 3, 1, 1 p
Fact sheets	http://www.orthoanswer.org/knee-leg/bakers-cyst/index.html





