

OUTPATIENT GP REFERRAL GUIDELINES UPPER LIMB

Southern Adelaide Local Health Network (SALHN)

Shoulder Osteoarthritis

Defined as

- Degeneration of articular cartilage causing pain, swelling and reduced movement
- Usually > 60 years
- Slow increase in pain
- Crepitus
- Loss of active + passive range of movement

Information Required for Referral

- History
- Onset, progression, duration, severity, any initiating factors
- · Range or movement of joint
- Functional limitations
- Associated medical conditions
- Any previous surgeries
- Any previous treatments
- Current medications

Investigations Required for Referral

- Xray AP/lateral/Axillary shoulder viewS
- Ultrasound check for rotator cuff rupture

Fax to

Orthopaedics Outpatients Upper Limb Clinic

Suggested GP Management

- NSAID
- Home Exercises
- Steroid Injection (IA)
- Refer if no improvement after conservative treatment > 3 months

Clinical Resources

08 83742591

Fax:

 American Academy of Orthopaedic Surgery. The treatment of Glenohumeral joint osteoarthritis Guidelines and Evidence Report V1 Dated 5.12.09

General Information to assist with referrals and the and Referral templates for FMC and RGH are available to download from the SALHN Outpatient Services website www.sahealth.sa.gov.au/SALHNoutpatients



