What to do if someone has taken your Methadone or Buprenorphine?

Administer naloxone, if available.

Ring **000** immediately if a child has swallowed Methadone or Buprenorphine. Give the paramedics the medication packaging.

If a child or an opiate naïve person may have taken your Methadone or Buprenorphine they may:

- > be sleepy or drowsy
- > be pale
- > be flat and have no energy
- perspire (sweat)
- > have trouble breathing

Naloxone is a life-saving medication that reverses the effects of opioid overdose. Ask your pharmacist for information on naloxone.

Helpful numbers

Ambulance attendance ph: **000** Poison Information Centre ph: 13 11 26 Not sure if someone has taken your Methadone or Buprenorphine. If you think a child or an opiate naïve person may have taken your Methadone or Buprenorphine:

- take them immediately to the emergency department of a hospital or call an ambulance
- > take the container with you
- > tell the staff the time you think they took the Methadone or Buprenorphine

Methadone, Buprenorphine (Subutex®, Suboxone®) and other drugs of dependence can be very dangerous to children and other people.

11 March 2020 DASSA:00443





Drug and Alcohol Services South Australia www.sahealth.sa.gov.au/dassa

Methadone

Buprenorphine

NOT FOR KIDS

DOs








DON'T'S...






DON'T ever tamper with the packaging.



DON'T'S...





DON'T keep **your** Methadone or Buprenorphine on you ...eg in handbag or pockets. Leave it locked up at home.

to anyone else.



