



Drug use in Adelaide Monitored by Wastewater Analysis

Project commissioned by Drug and Alcohol Services
South Australia (DASSA)

Analyses performed by:
School of Pharmacy and Medical Sciences
University of South Australia

August 2017

Please note that drug consumption levels may vary slightly from report to report due to adjustments made to sewerage flow rates in some of the treatment plants.



Purpose of the project

- > To determine the prevalence of drug use in South Australia, initially in metropolitan Adelaide, through wastewater analysis.



Wastewater analysis CAN tell us:

- > The pattern of drug use over the week.
- > The prevalence of drug use every two months from December 2011.



Wastewater analysis **CANNOT** tell us:

- > The characteristics of drug users.
- > In what regions of metropolitan Adelaide drug consumption is occurring.
- > The form and way drugs were taken.




Wastewater sampling

- > Sampling every two months from Adelaide Metropolitan wastewater, commenced in December 2011.

- > Drugs tested:
 - Stimulants: cocaine, MDMA, and methamphetamine.
 - Opioids: morphine, codeine, methadone, oxycodone, fentanyl and heroin.
 - Cannabis (THC).
 - Nicotine.
 - Anabasine (a tobacco-specific alkaloid).
 - Testing for alcohol has commenced, and results will appear in future iterations of this report.

Stimulants


Methamphetamine consumption increased from 2012-2016, with decreases in the first half of 2017 followed by an increase in August.



Average consumption (dose/week/1000 people) of methamphetamine for 2012-2015. Weekly consumption (dose/week/1000 people) bi-monthly from February 2016.

Stimulants


Methamphetamine consumption increases slightly on weekends.



Average consumption (dose/day/1000 people) of methamphetamine over the week.
Dose = 30 mg.

Stimulants


Cocaine consumption has increased since 2015.



Average consumption (dose/day/1000 people) of cocaine for 2012-2015. Weekly consumption (dose/week/1000 people bi-monthly from February 2016. Dose = 100 mg.

Stimulants


Cocaine consumption is higher on weekends.



Average consumption (dose/day/1000 people) of cocaine over the week. Dose = 100 mg.

Stimulants


Ecstasy (MDMA) consumption decreased from 2012 to 2016 with a small increase in 2017.



Average consumption (dose/day/1000 people) of MDMA for 2012-2015. Weekly consumption (dose/week/1000 people bi-monthly from February 2016. Dose = 100 mg.


Stimulants

Ecstasy (MDMA) consumption is higher on weekends.



Average consumption (dose/day/1000 people) of MDMA over the week. Dose = 100 mg.

Stimulants



Average consumption (dose/day/1000 people) 2012-2015 . Weekly consumption (dose/week/1000 people) of cocaine (100mg dose), MDMA (100mg dose) and methamphetamine (30 mg dose) from Feb 2016.



Stimulants - summary

- > **Methamphetamine:**
 - > Highest level of consumption of the stimulants.
 - > Steady increase in levels of consumption since December 2012; decreases in the first half of 2017 were followed by an increase in August.
- > **Cocaine:**
 - > Consumption has increased since 2015 but overall consumption is low.
- > **Ecstasy (MDMA):**
 - > Consumption decreased from 2012 to 2016 with a small increase in 2017.




Opioids

- > Opioids are a class of drugs that are used for pain relief (e.g. codeine, morphine) or for the treatment of opioid dependence (e.g. methadone).
- > Codeine in the samples can originate from prescription or over the counter medications.
- > Morphine, methadone, oxycodone and fentanyl can be used legally on prescription or may be sourced illegally.
- > The analysis of opioids, except for heroin, cannot differentiate illicit from licit use.

Opioids


Heroin consumption has decreased since 2013.



Average consumption (dose/week/1000 people) of heroin for 2013-2015. Weekly consumption (dose/week/1000 people) bi-monthly from February 2016. Dose for calculation = 20 mg.


Opioids

Heroin consumption is constant over the week.



Average consumption (dose/day/1000 people) of heroin over the week. Dose = 20mg.


Medicinal Opioids



Daily consumption (dose/day/1000 people) of codeine (200mg dose), morphine, methadone (100mg dose), oxycodone (10mg dose) and fentanyl (0.2mg dose) from Dec 2011.

Opioids


Codeine consumption is constant over the week.



Average consumption (dose/day/1000 people) of codeine over the week. Dose = 200 mg.

Opioids


Methadone consumption is low and constant over the week.



Average consumption (dose/day/1000 people) of methadone over the week. Dose = 100 mg.

Opioids


Morphine consumption is constant over the week.



Average daily consumption (dose/day/1000 people) of morphine over the week.

Opioids


Oxycodone consumption is constant over the week.



Average consumption (dose/day/1000 people) of oxycodone over the week. Dose = 10 mg.

Opioids

Fentanyl consumption is constant over the week.



Average consumption (dose/day/1000 people) of fentanyl over the week. Dose = 0.2mg.




Opioids - summary

- > Oxycodone and fentanyl showed increases in consumption over the period of data collection, but have decreased in 2016 and 2017 to levels seen in 2012 and 2013.
- > Heroin consumption has decreased since 2013 and has stayed relatively low.

Cannabis


There has been a small downward trend in cannabis consumption levels since 2012.



Average consumption (dose/week/1000 people) of THC for 2012-2015. Weekly consumption (dose/week/1000 people) bi-monthly from February 2016. Dose = 125 mg.

Cannabis


Cannabis consumption is fairly constant over the week.



Average consumption (dose/day/1000 people) of THC over the week. Dose = 125 mg.

Nicotine*

Nicotine consumption has shown a gradual decline since 2012.




Average consumption (dose/day/1000 people) of nicotine for 2012-2015. Weekly consumption (dose/week/1000 people) bi-monthly from February 2016. Dose = 1mg.

*Does not differentiate between tobacco and nicotine replacement therapy (NRT) use

Nicotine


Nicotine consumption is fairly constant over the week.



Average consumption (dose/day/1000 people) of nicotine over the week. Dose = 1mg.

Anabesine*

Anabesine has shown a decline since December 2015.



Average weekly excretion (mg excreted/week/ 1000 people) from December 2015. As yet there is no excretion rate to convert to number of cigarettes smoked.

*Tobacco specific alkaloid



Summary

- > Methamphetamine is the predominant stimulant consumed in metropolitan Adelaide and has shown a steady increase in consumption since 2012. Decreases in the first half of 2017 were followed by an increase in August.
- > Other stimulants are consumed at much lower rates.
- > Stimulant consumption increases on the weekends.
- > Opioid consumption is relatively constant across the week.



Summary (continued)

- > Oxycodone and fentanyl consumption increased to 2015, and has decreased in 2016 and 2017.
- > Cannabis consumption has shown a small downward trend over the sampling period and it is used constantly over the week.
- > Nicotine consumption has shown a very gradual decline over the sampling period and is fairly constant over the week.
- > Anabesine has also shown a decline over the reporting period.