

OUTPATIENT GP REFERRAL GUIDELINES GASTROENTEROLOGY Southern Adelaide Local Health Network (SALHN)

Nausea and Vomiting

Chronic nausea/vomiting is defined by symptoms >1 month duration

Information Required

- Presence of Red flags
- Duration, frequency and severity
- Smoking, alcohol, medications (including recreational drugs)

Investigations Required

- FBE, EUC, fasting glucose, LFTs, CRP, TFTs
- Urinalysis and urine HCG if premenopausal female
- Hepatobiliary abdominal US and abdominal X-Ray

Fax Referrals to

 Gastroenterology Outpatient Clinic Flinders Medical Centre 8204 5555

Red Flags

Dysphagia

Jyspriagia

Haematemesis
Faeculant vomiting

Persistent abdominal pain

Weight loss

Suggested GP Management

- Exclude non-gastrointestinal causes Intracranial pathology, labyrinthitis, pregnancy, thyroid disease, medications
- Advise small, frequent, low fat meals

Acute nausea and vomiting

- Non toxic
 - Maintain oral hydration, antiemetics
- Febrile, dehydrated
 - Perform basic blood and urine investigations as below
 - Consider erect, supine abdominal X-ray
 - Consider referral to Emergency Department if unable to tolerate oral fluids

Chronic nausea and vomiting

- Refer if age ≥50yrs or Red flags
- Trial proton pump inhibitor or antiemetics; refer if no response after 4 weeks

Clinical Resources

- American Gastroenterological Association Medical Position Statement: Nausea and Vomiting Gastroenterology 2001;120:261-62.
- Therapeutic Guidelines www.tg.org.au

General Information to assist with referrals and the and Referral templates for FMC are available to download from the SAHN Outpatient Services website www.sahealth.sa.gov.au/SALHNoutpatients



Version	Date from	Date to	Amendment
2.1	Nov 2017	Nov 2019	Removed all RGH details